

08-30-00

A

UTILITY PATENT APPLICATION TRANSMITTAL

(for new applications under 37 C.F.R. § 1.53(b))

08/29/00
JC821 U.S. PROJC864 U.S. PRO
09/650134
08/29/00

Invention Docket No.: C4007(C)
 Applicant: Ronald HAGE; Simon Marinus VEERMAN
 Title: COMPOSITION AND METHOD FOR BLEACHING A SUBSTRATE
 Express Mail Label No.: EK 942 558 917 US
 Date Deposited: August 29, 2000
 UNUS No.: Y2-0187-UNI

To: Assistant Commissioner for Patents
 Box: Patent Application
 Washington, D.C. 20231

APPLICATION ELEMENTS

1. ☒ Fee Calculation (Box 13) and Authorization (Triplicate copies of this form are enclosed)
2. ☒ Specification and Claims (97) Total Pages
3. ☐ Formal or Informal Drawings () Total Sheets
4. ☒ Unexecuted Declaration
- ☐ Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - a. ☐ Computer Readable Copy
 - b. ☐ Paper copy (identical to computer copy)
 - c. ☐ Statement verifying identify of above copies.

ACCOMPANYING APPLICATION PARTS

6. ☐ Information Disclosure Statement (IDS)/PTO-1449
7. ☐ Copies of IDS citations
8. ☐ Preliminary Amendment
9. ☒ Two (2) Return Receipt Postcards
10. ☐ Certified Copy of Priority Document
11. ☒ The benefit under 35 U.S.C. § 119 is claimed of the filing of PCT Application No. PCT/GB99/02876 filed September 1, 1999; PCT Application No. PCT/GB99/02878 filed September 1, 1999; and United Kingdom Application No. 0004990.8 filed March 1, 2000.
12. ☐ Other:
13. ☒ The FILING FEE (including any claims introduced or cancelled by Preliminary Amendment) is calculated below:

CLAIMS				
FOR	NUMBER FILED	NUMBER EXTRA	RATE	BASIC FEE \$690.00
Total Claims	21 - 20	1	X \$ 18.00	\$ 18.00
Independent Claims	4 - 3	1	X \$ 78.00	\$ 78.00
Multiple Claims	<u>Yes</u> <u>No</u>		X \$260.00	
TOTAL FILING FEE . . .				\$786.00

14. ☒ Charge \$786.00 to Deposit Acct. #12-1155. Triplicate copies of this letter are enclosed.
15. ☒ The Commissioner is hereby authorized to charge any additional fees, which may be required, including all required fees under
- ☒ 37 C.F.R. § 1.16;
 - ☒ 37 C.F.R. § 1.17;
 - ☒ 37 C.F.R. § 1.18.
16. ☒ **Correspondence Address:**



00201

PATENT TRADEMARK OFFICE

Respectfully submitted,

Milton L. Honig
Attorney of Record
Reg. #28,617

MLH/mt
(201) 840-2403

COMPOSITION AND METHOD FOR BLEACHING A SUBSTRATEFIELD OF INVENTION

5 This invention relates to compositions and methods for catalytically bleaching substrates with atmospheric oxygen, using a metal-ligand complex as catalyst, which compositions are formulated as liquids. This invention also relates to a method of treating textiles, such as laundry fabrics, using
10 a metal-ligand complex as catalyst whereby bleaching with atmospheric oxygen is catalysed after the treatment, wherein the treatment composition is formulated as a liquid.

BACKGROUND OF INVENTION

15 Peroxygen bleaches are well known for their ability to remove stains from substrates. Traditionally, the substrate is subjected to hydrogen peroxide, or to substances which can generate hydroperoxyl radicals, such as inorganic or
20 organic peroxides. Generally, these systems must be activated. One method of activation is to employ wash temperatures of 60°C or higher. However, these high temperatures often lead to inefficient cleaning, and can also cause premature damage to the substrate.

25 A preferred approach to generating hydroperoxyl bleach species is the use of inorganic peroxides coupled with organic precursor compounds. These systems are employed for many commercial laundry powders. For example, various
30 European systems are based on tetraacetyl ethylenediamine (TAED) as the organic precursor coupled with sodium

perborate or sodium percarbonate, whereas in the United States laundry bleach products are typically based on sodium nonanoyloxybenzenesulphonate (SNOBS) as the organic precursor coupled with sodium perborate.

5

Precursor systems are generally effective but still exhibit several disadvantages. For example, organic precursors are moderately sophisticated molecules requiring multi-step manufacturing processes resulting in high capital costs.

10 Also, precursor systems have large formulation space requirements so that a significant proportion of a laundry powder must be devoted to the bleach components, leaving less room for other active ingredients and complicating the development of concentrated powders. Moreover, precursor
15 systems do not bleach very efficiently in countries where consumers have wash habits entailing low dosage, short wash times, cold temperatures and low wash liquor to substrate ratios.

20 Alternatively, or additionally, hydrogen peroxide and peroxy systems can be activated by bleach catalysts, such as by complexes of iron and the ligand N4Py (*i.e.* N, N-bis(pyridin-2-yl-methyl)-bis(pyridin-2-yl)methylamine) disclosed in WO95/34628, or the ligand Tpen (*i.e.* N, N, N',
25 N'-tetra(pyridin-2-yl-methyl)ethylenediamine) disclosed in WO97/48787. These publications do not foresee a role in providing storage stable liquid bleaching compositions even if, according to these publications, molecular oxygen may be used as the oxidant as an alternative to peroxide generating
30 systems.

As discussed by N.J. Milne in J. of Surfactants and Detergents, Vol 1, no 2, 253-261 (1998), it has long been thought desirable to be able to use atmospheric oxygen (air) as the source for a bleaching species. The use of atmospheric oxygen (air) as the source for a bleaching species would avoid the need for costly hydroperoxyl generating systems. Unfortunately, air as such is kinetically inert towards bleaching substrates and exhibits no bleaching ability. Recently some progress has been made in this area. For example, WO 97/38074 reports the use of air for oxidising stains on fabrics by bubbling air through an aqueous solution containing an aldehyde and a radical initiator. A broad range of aliphatic, aromatic and heterocyclic aldehydes is reported to be useful, particularly para-substituted aldehydes such as 4-methyl-, 4-ethyl- and 4-isopropyl benzaldehyde, whereas the range of initiators disclosed includes N-hydroxysuccinimide, various peroxides and transition metal coordination complexes.

However, although this system employs molecular oxygen from the air, the aldehyde component and radical initiators such as peroxides are consumed during the bleaching process. These components must therefore be included in the composition in relatively high amounts so as not to become depleted before completion of the bleaching process in the wash cycle. Moreover, the spent components represent a waste of resources as they can no longer participate in the bleaching process.

Accordingly, it would be desirable to be able to provide a bleaching system based on atmospheric oxygen or air that

does not rely primarily on hydrogen peroxide or a hydroperoxyl generating system, and that does not require the presence of organic components such as aldehydes that are consumed in the process. Moreover, it would be
5 desirable to provide such a bleaching system that is effective in aqueous medium.

It may also be noted that the known art teaches a bleaching effect only as long as the substrate is being subjected to
10 the bleaching treatment. Thus, there is no expectation that hydrogen peroxide or peroxy bleach systems could continue to provide a bleaching effect on a treated substrate, such as a laundry fabric after washing and drying, since the bleaching species themselves or any activators necessary for the
15 bleaching systems would be assumed to be removed from the substrate, or consumed or deactivated, on completing the wash cycle and drying.

It would be therefore also be desirable to be able to treat
20 a textile such that, after the treatment is completed, a bleaching effect is observed on the textile. Furthermore, it would be desirable to be able to provide a bleach treatment for textiles such as laundry fabrics whereby residual bleaching occurs in the presence of air when the
25 treated fabric has been treated and is dry. It would be desirable for the residual bleaching of dry textiles to be conducted irrespective of exposure to light.

A further disadvantage associated with conventional
30 bleaching compositions based on hydrogen peroxide or peroxy systems such those containing organic peroxyacids is that

the compositions tend to be chemically or physically unstable in the presence of liquid solvents, carriers or other liquid components such as surfactants, particularly when formulated as aqueous compositions. Consequently, when formulated as liquids, these bleaching compositions on the one hand do not exhibit satisfactory storage stability, resulting in a rapid loss of bleaching activity or in a loss of structural integrity, for example phase separation, or require the incorporation of additional stabilising systems to minimise these effects with attendant disadvantages in terms of cost or processing. Decomposition of a hydrogen peroxide or peroxy liquid bleaching composition in a sealed container leads to an increase in the internal pressure of the sealed container. The increase in the internal pressure leads to the possibility of the sealed container rupturing in a dangerous manner. In the presence of surfactants, decomposition of the hydrogen peroxide or peroxy liquid bleaching composition leads to foaming of the composition. On the other hand, liquid bleaching compositions are conveniently dosed into containers for storage or for use, or otherwise handled, and are desired by the consumer, particularly in the United States of America.

It would therefore also be desirable to be able to provide a bleaching composition in the form of a liquid, which is chemically and physically stable, without at least some of the disadvantages hitherto associated with liquid bleaching compositions. It would furthermore be desirable to be able to provide chemically and physically storage stable detergent bleaching compositions or rinse conditioning bleach compositions in the form of a liquid.

Application W000/29537, filed 9 November 1999, was published after the filing date of the present application disclosing theoretical examples of compositions for bleaching with a transition metal complex in the absence of an added
5 peroxygen bleach. Application W000/29537 has no evidence of efficacy and includes two classes of ligands: some cross-bridged macrocyclic ligands and some macrocyclic ligands. The macrocyclic ligands are disclosed as manganese complexes and are not found in the priority document of W000/29537;
10 namely US serial number 60/108,292 filed 13 November 1998. The theoretical examples given are for a heavy-duty granular laundry detergent and heavy-duty liquid laundry detergent. In both these examples the exemplified bleach catalyst is 5,12-dimethyl-1,5,8,12-tetra-bicyclo[6.6.2.]hexadecane
15 manganese (II) chloride. There are no examples demonstrating any bleaching effect. The use of manganese complexes in laundry applications is less preferred because of dye/textile damage under specific conditions.

20 SUMMARY OF INVENTION

We have now found that it is possible to achieve a chemically and physically stable bleaching composition in the form of a liquid, by using an organic substance that
25 forms a complex which catalyses the bleaching of substrates using atmospheric oxygen or air, and formulating the organic substance in a liquid that is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system. Moreover, we have found that these organic
30 substances can be formulated together with detergent or rinse conditioning agents, in a liquid that is substantially

devoid of peroxygen bleach or a peroxy-based or -generating bleach system, to provide chemically and physically stable detergent bleaching compositions or rinse conditioning bleach compositions, in the form of a liquid.

5

Accordingly, in a first aspect, the present invention a liquid bleaching composition comprising an organic substance which forms a complex with a transition metal, the complex catalysing bleaching of a substrate by atmospheric oxygen, and a liquid carrier or solvent, wherein the composition is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system. The composition is therefore preferably insensitive or stable to catalase, which acts on peroxy species.

15

In a second aspect, the present invention provides a method of bleaching a substrate comprising applying to the substrate a liquid bleaching composition that comprises an organic substance which forms a complex with a transition metal, the complex catalysing bleaching of the substrate by atmospheric oxygen, and a liquid carrier or solvent, wherein the composition is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system.

25

Furthermore, in a third aspect, the present invention provides the use of an organic substance which forms a complex with a transition metal, the complex catalysing bleaching of a substrate by the atmospheric oxygen, as a catalytic bleaching agent in a liquid bleaching composition substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system.

30

We have also found that the liquid bleaching compositions in accordance with the present invention are surprisingly effective in catalysing bleaching of substrates by atmospheric oxygen after treatment of the substrate.

5

Accordingly, in a fourth aspect, the present invention provides a method of treating a textile by contacting the textile with a liquid bleaching composition that comprises an organic substance which forms a complex with a transition metal, the complex catalysing bleaching by atmospheric oxygen, and a liquid carrier or solvent, wherein the composition is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system, whereby the complex catalyses bleaching of the textile by atmospheric oxygen after the treatment.

The present invention requires all or the majority of the bleaching species in the liquid bleaching composition (on an equivalent weight basis) to be derived from atmospheric oxygen. Thus, the liquid composition will be made wholly or substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system. The organic substance is a catalyst for the bleaching process and, as such, is not consumed but can continue to participate in the bleaching process. Since the bleaching system of the type used in the liquid bleaching composition is catalytically activated and the bleaching species is derived from atmospheric oxygen, the present invention is advantageous in that it provides a bleaching composition which is not only convenient to handle by virtue of being in the form of a liquid, but which also is both cost-effective and environmentally friendly.

The liquid bleaching composition may be formulated as a concentrated bleaching liquid for direct application to a substrate, or for application to a substrate following dilution, such as dilution before or during use of the liquid composition by the consumer or in washing apparatus.

The liquid bleaching composition can for example be formulated as an aqueous medium, or so as to be dispersable into an aqueous medium, and is therefore particularly applicable to bleaching of laundry fabrics. Therefore, whilst the composition and method according to the present invention may be used for bleaching any suitable substrate, the preferred substrate is a laundry fabric. Bleaching may be carried out by simply leaving the substrate in contact for a sufficient period of time with a bleach medium constituted by or prepared from the liquid bleaching composition. Preferably, however, the bleach medium on or containing the substrate is agitated.

An advantage of the method according to the fourth aspect of the invention is that, by enabling a bleaching effect even after the textile has been treated, the benefits of bleaching can be prolonged on the textile. Furthermore, since a bleaching effect is conferred to the textile after the treatment, the treatment itself, such as a laundry wash cycle, may for example be shortened.

The present invention also extends to a commercial package comprising a liquid bleaching composition comprising a ligand or complex as defined below together with instructions for its use.


The present invention also extends to use of a ligand or complex as defined below in the manufacture of a liquid bleaching composition, the bleaching composition substantially devoid of peroxygen bleach or a peroxy-based or peroxy-generating bleach system.

DETAILED DESCRIPTION OF THE INVENTION

The catalyst may comprise a preformed complex of a ligand and a transition metal. Alternatively, the catalyst may comprise a free ligand that complexes with a transition metal already present in the water or that complexes with a transition metal present in the substrate. The catalyst may also be included in the form of a composition of a free ligand or a transition metal-substitutable metal-ligand complex, and a source of transition metal, whereby the complex is formed *in situ* in the medium. It is preferred that the catalyst is a pentadentate ligand or complex thereof.

The ligand forms a complex with one or more transition metals, in the latter case for example as a dinuclear complex. Suitable transition metals include for example: manganese in oxidation states II-V, iron II-V, copper I-III, cobalt I-III, titanium II-IV, tungsten IV-VI, vanadium II-V and molybdenum II-VI.

The transition metal complex preferably is of the general formula:



in which:

M represents a metal selected from Mn(II)-(III)-(IV)-(V), Cu(I)-(II)-(III), Fe (II)-(III)-(IV)-(V), Co(I)-(II)-(III), Ti(II)-(III)-(IV), V(II)-(III)-(IV)-(V), Mo(II)-(III)-(IV)-(V)-(VI) and W(IV)-(V)-(VI), preferably from Fe(II)-(III)-(IV)-(V);

L represents the ligand, preferably N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, or its protonated or deprotonated analogue;


X represents a coordinating species selected from any mono, bi or tri charged anions and any neutral molecules able to coordinate the metal in a mono, bi or tridentate manner;

Y represents any non-coordinated counter ion;
a represents an integer from 1 to 10;
k represents an integer from 1 to 10;
n represents zero or an integer from 1 to 10;
m represents zero or an integer from 1 to 20.

Preferably, the complex is an iron complex comprising the ligand N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane. However, it will be appreciated that the pretreatment method of the present invention may instead, or additionally, use other ligands and transition metal complexes, provided that the complex formed is capable of catalysing stain bleaching by atmospheric oxygen. Suitable classes of ligands are described below:

(A) Ligands of the general formula (IA):

- 12 -




(IA)

wherein

5 Z1 groups independently represent a coordinating group selected from hydroxy, amino, -NHR or -N(R)₂ (wherein R=C₁₋₆-alkyl), carboxylate, amido, -NH-C(NH)NH₂, hydroxyphenyl, a heterocyclic ring optionally substituted by one or more functional groups E or a heteroaromatic ring optionally substituted by one or more functional groups E, the heteroaromatic ring being selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole;

15 Q1 and Q3 independently represent a group of the formula:



20 wherein

$5 \geq a+b+c \geq 1$; $a=0-5$; $b=0-5$; $c=0-5$; $n=0$ or 1
(preferably $n=0$);

Y independently represents a group selected from -O-, -
5 S-, -SO-, -SO₂-, -C(O)-, arylene, alkylene, heteroarylene,
heterocycloalkylene, -(G)P-, -P(O)- and -(G)N-, wherein G
is selected from hydrogen, alkyl, aryl, arylalkyl,
cycloalkyl, each except hydrogen being optionally
substituted by one or more functional groups E;




10 R5, R6, R7, R8 independently represent a group selected
from hydrogen, hydroxyl, halogen, -R and -OR, wherein R
represents alkyl, alkenyl, cycloalkyl, heterocycloalkyl,
aryl, heteroaryl or a carbonyl derivative group, R being
15 optionally substituted by one or more functional groups E,
or R5 together with R6, or R7 together with R8, or
both, represent oxygen,

or R5 together with R7 and/or independently R6 together
with R8, or R5 together with R8 and/or independently R6
20 together with R7, represent C₁₋₆-alkylene optionally
substituted by C₁₋₄-alkyl, -F, -Cl, -Br or -I;

T represents a non-coordinated group selected from
hydrogen, hydroxyl, halogen, -R and -OR, wherein R
represents alkyl, alkenyl, cycloalkyl, heterocycloalkyl,
25 aryl, arylalkyl, heteroaryl or a carbonyl derivative group,
R being optionally substituted by one or more functional
groups E (preferably T= -H, -OH, methyl, methoxy or benzyl);

U represents either a non-coordinated group T
30 independently defined as above or a coordinating group of
the general formula (IIA), (IIIA) or (IVA):

- 14 -



wherein


Q2 and Q4 are independently defined as for Q1 and Q3;

Q represents -N(T)- (wherein T is independently defined as above), or an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole;

Z2 is independently defined as for Z1;

Z3 groups independently represent -N(T)- (wherein T is independently defined as above);

Z4 represents a coordinating or non-coordinating group
 5 selected from hydrogen, hydroxyl, halogen, -NH-C(NH)NH₂, -R and -OR, wherein R= alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or a carbonyl derivative group, R being optionally substituted by one or more functional groups E, or Z4 represents a group of the general
 10 formula (IIAa):



and

15 $1 \leq j < 4$.

Preferably, Z1, Z2 and Z4 independently represent an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine,
 20 pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole. More preferably, Z1, Z2 and Z4 independently represent groups selected from optionally substituted pyridin-2-yl, optionally substituted
 25 imidazol-2-yl, optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl. Most preferred is that Z1, Z2 and Z4 each represent optionally substituted pyridin-2-yl.

The groups Z1, Z2 and Z4 if substituted, are preferably substituted by a group selected from C₁₋₄-alkyl, aryl, arylalkyl, heteroaryl, methoxy, hydroxy, nitro, amino, carboxyl, halo, and carbonyl. Preferred is that Z1, Z2 and
5 Z4 are each substituted by a methyl group. Also, we prefer that the Z1 groups represent identical groups.

Each Q1 preferably represents a covalent bond or C1-C4-alkylene, more preferably a covalent bond, methylene or
10 ethylene, most preferably a covalent bond.


Group Q preferably represents a covalent bond or C1-C4-alkylene, more preferably a covalent bond.

15 The groups R5, R6, R7, R8 preferably independently represent a group selected from -H, hydroxy-C₀-C₂₀-alkyl, halo-C₀-C₂₀-alkyl, nitroso, formyl-C₀-C₂₀-alkyl, carboxyl-C₀-C₂₀-alkyl and esters and salts thereof, carbamoyl-C₀-C₂₀-alkyl, sulfo-C₀-C₂₀-alkyl and esters and salts thereof, sulfamoyl-C₀-C₂₀-
20 alkyl, amino-C₀-C₂₀-alkyl, aryl-C₀-C₂₀-alkyl, C₀-C₂₀-alkyl, alkoxy-C₀-C₈-alkyl, carbonyl-C₀-C₆-alkoxy, and C₀-C₂₀-alkylamide. Preferably, none of R5-R8 is linked together.

Non-coordinated group T preferably represents hydrogen,
25 hydroxy, methyl, ethyl, benzyl, or methoxy.

In one aspect, the group U in formula (IA) represents a coordinating group of the general formula (IIA):

- 17 -



According to this aspect, it is preferred that Z2 represents
 5 an optionally substituted heterocyclic ring or an optionally
 substituted heteroaromatic ring selected from pyridine,
 pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole,
 quinoline, quinoxaline, triazole, isoquinoline, carbazole,
 indole, isoindole, oxazole and thiazole, more preferably
 10 optionally substituted pyridin-2-yl or optionally
 substituted benzimidazol-2-yl.

It is also preferred, in this aspect, that Z4 represents an
 optionally substituted heterocyclic ring or an optionally
 15 substituted heteroaromatic ring selected from pyridine,
 pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole,
 quinoline, quinoxaline, triazole, isoquinoline, carbazole,
 indole, isoindole, oxazole and thiazole, more preferably
 optionally substituted pyridin-2-yl, or a non-coordinating
 20 group selected from hydrogen, hydroxy, alkoxy, alkyl,
 alkenyl, cycloalkyl, aryl, or benzyl.

In preferred embodiments of this aspect, the ligand is
 selected from:


25 1,1-bis(pyridin-2-yl)-N-methyl-N-(pyridin-2-
 ylmethyl)methylamine;
 1,1-bis(pyridin-2-yl)-N,N-bis(6-methyl-pyridin-2-
 ylmethyl)methylamine;

1,1-bis(pyridin-2-yl)-N,N-bis(5-carboxymethyl-pyridin-2-ylmethyl)methylamine;


1,1-bis(pyridin-2-yl)-1-benzyl-N,N-bis(pyridin-2-ylmethyl)methylamine; and

5 1,1-bis(pyridin-2yl)-N,N-bis(benzimidazol-2-ylmethyl)methylamine.

In a variant of this aspect, the group Z4 in formula (IIA) represents a group of the general formula (IIAa):




In this variant, Q4 preferably represents optionally substituted alkylene, preferably -CH₂-CHOH-CH₂- or -CH₂-CH₂-CH₂-. In a preferred embodiment of this variant, the ligand is:



wherein -Py represents pyridin-2-yl.

In another aspect, the group U in formula (IA) represents a coordinating group of the general formula (IIIA):

- 19 -




(IIIA)

wherein j is 1 or 2, preferably 1.

5

According to this aspect, each Q2 preferably represents $(\text{CH}_2)_n$ ($n=2-4$), and each Z3 preferably represents $-\text{N}(\text{R})-$ wherein R = $-\text{H}$ or C_{1-4} -alkyl, preferably methyl.


10 In preferred embodiments of this aspect, the ligand is selected from:



wherein $-\text{Py}$ represents pyridin-2-yl.

15

In yet another aspect, the group U in formula (IA) represents a coordinating group of the general formula (IVA):




(IVA)

20


In this aspect, Q preferably represents -N(T)- (wherein T= -H, methyl, or benzyl) or pyridin-diyl.

In preferred embodiments of this aspect, the ligand is
5 selected from:



wherein -Py represents pyridin-2-yl, and -Q- represents
10 pyridin-2,6-diyl.

(B) Ligands of the general formula (IB):




(IB)

wherein

20 $n = 1$ or 2 , whereby if $n = 2$, then each $-\text{Q}_3-\text{R}_3$ group is independently defined;

R_1, R_2, R_3, R_4 independently represent a group selected from hydrogen, hydroxyl, halogen, $-\text{NH}-\text{C}(\text{NH})\text{NH}_2$, $-\text{R}$ and $-\text{OR}$, wherein $\text{R} =$ alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or a carbonyl derivative group, R being
 5 optionally substituted by one or more functional groups E ,

Q_1, Q_2, Q_3, Q_4 and Q independently represent a group of the formula:



wherein

$5 \geq a+b+c \geq 1$; $a=0-5$; $b=0-5$; $c=0-5$; $n=1$ or 2 ;

15 Y independently represents a group selected from $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})-$, arylene, alkylene, heteroarylene, heterocycloalkylene, $-(\text{G})\text{P}-$, $-\text{P}(\text{O})-$ and $-(\text{G})\text{N}-$, wherein G is selected from hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, each except hydrogen being optionally
 20 substituted by one or more functional groups E ;

$\text{R5}, \text{R6}, \text{R7}, \text{R8}$ independently represent a group selected from hydrogen, hydroxyl, halogen, $-\text{R}$ and $-\text{OR}$, wherein R represents alkyl, alkenyl, cycloalkyl, heterocycloalkyl,

aryl, heteroaryl or a carbonyl derivative group, R being optionally substituted by one or more functional groups E, or R5 together with R6, or R7 together with R8, or both, represent oxygen,

5 or R5 together with R7 and/or independently R6 together with R8, or R5 together with R8 and/or independently R6 together with R7, represent C₁₋₆-alkylene optionally substituted by C₁₋₄-alkyl, -F, -Cl, -Br or -I,



10 provided that at least two of R₁, R₂, R₃, R₄ comprise coordinating heteroatoms and no more than six heteroatoms are coordinated to the same transition metal atom.

At least two, and preferably at least three, of R₁, R₂, R₃, R₄ independently represent a group selected from carboxylate, amido, -NH-C(NH)NH₂, hydroxyphenyl, an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole.

20 Preferably, substituents for groups R₁, R₂, R₃, R₄, when representing a heterocyclic or heteroaromatic ring, are selected from C₁₋₄-alkyl, aryl, arylalkyl, heteroaryl, methoxy, hydroxy, nitro, amino, carboxyl, halo, and carbonyl.

The groups Q₁, Q₂, Q₃, Q₄ preferably independently represent a group selected from -CH₂- and -CH₂CH₂-.

Group Q is preferably a group selected from $-(CH_2)_{2-4}-$, $-CH_2CH(OH)CH_2-$,




5 wherein R represents $-H$ or C_{1-4} -alkyl.

Preferably, Q_1 , Q_2 , Q_3 , Q_4 are defined such that $a=b=0$, $c=1$ and $n=1$, and Q is defined such that $a=b=0$, $c=2$ and $n=1$.

10 The groups R_5 , R_6 , R_7 , R_8 preferably independently represent a group selected from $-H$, hydroxy- C_0 - C_{20} -alkyl, halo- C_0 - C_{20} -alkyl, nitroso, formyl- C_0 - C_{20} -alkyl, carboxyl- C_0 - C_{20} -alkyl and esters and salts thereof, carbamoyl- C_0 - C_{20} -alkyl, sulfo- C_0 - C_{20} -alkyl and esters and salts thereof, sulfamoyl- C_0 - C_{20} -alkyl, amino- C_0 - C_{20} -alkyl, aryl- C_0 - C_{20} -alkyl, C_0 - C_{20} -alkyl, alkoxy- C_0 - C_8 -alkyl, carbonyl- C_0 - C_6 -alkoxy, and C_0 - C_{20} -alkylamide. Preferably, none of R_5 - R_8 is linked together.

In a preferred aspect, the ligand is of the general formula

20 (IIB):



(IIB)

25

wherein

Q_1, Q_2, Q_3, Q_4 are defined such that $a=b=0$, $c=1$ or 2 and $n=1$;

Q is defined such that $a=b=0$, $c=2,3$ or 4 and $n=1$; and

$R_1, R_2, R_3, R_4, R_7, R_8$ are independently defined as for
5 formula (I).

Preferred classes of ligands according to this aspect, as represented by formula (IIB) above, are as follows:

10 (i) ligands of the general formula (IIB) wherein:

R_1, R_2, R_3, R_4 each independently represent a coordinating group selected from carboxylate, amido, $-NH-C(NH)NH_2$, hydroxyphenyl, an optionally substituted heterocyclic ring or an optionally substituted
15 heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole.

20 In this class, we prefer that:

Q is defined such that $a=b=0$, $c=2$ or 3 and $n=1$;

R_1, R_2, R_3, R_4 each independently represent a coordinating group selected from optionally substituted pyridin-2-yl, optionally substituted imidazol-2-yl,
25 optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl.

(ii) ligands of the general formula (IIB) wherein:

R_1, R_2, R_3 each independently represent a coordinating
30 group selected from carboxylate, amido, $-NH-C(NH)NH_2$, hydroxyphenyl, an optionally substituted heterocyclic ring

or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole; and

R_4 represents a group selected from hydrogen, C_{1-20} optionally substituted alkyl, C_{1-20} optionally substituted arylalkyl, aryl, and C_{1-20} optionally substituted NR_3^+ (wherein $R=C_{1-8}$ -alkyl).

In this class, we prefer that:

Q is defined such that $a=b=0$, $c=2$ or 3 and $n=1$;

R_1 , R_2 , R_3 each independently represent a coordinating group selected from optionally substituted pyridin-2-yl, optionally substituted imidazol-2-yl, optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl; and

R_4 represents a group selected from hydrogen, C_{1-10} optionally substituted alkyl, C_{1-5} -furanyl, C_{1-5} optionally substituted benzylalkyl, benzyl, C_{1-5} optionally substituted alkoxy, and C_{1-20} optionally substituted N^+Me_3 .

(iii) ligands of the general formula (IIB) wherein:

R_1 , R_4 each independently represent a coordinating group selected from carboxylate, amido, $-NH-C(NH)NH_2$, hydroxyphenyl, an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole; and

R_2 , R_3 each independently represent a group selected from hydrogen, C_{1-20} optionally substituted alkyl, C_{1-20} optionally substituted arylalkyl, aryl, and C_{1-20} optionally substituted NR_3^+ (wherein $R=C_{1-8}$ -alkyl).

5

In this class, we prefer that:

Q is defined such that $a=b=0$, $c=2$ or 3 and $n=1$;

R_1 , R_4 each independently represent a coordinating group selected from optionally substituted pyridin-2-yl, optionally substituted imidazol-2-yl, optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl; and

R_2 , R_3 each independently represent a group selected from hydrogen, C_{1-10} optionally substituted alkyl, C_{1-5} -furanyl, C_{1-5} optionally substituted benzylalkyl, benzyl, C_{1-5} optionally substituted alkoxy, and C_{1-20} optionally substituted N^+Me_3 .

Examples of preferred ligands in their simplest forms are:

20

N,N',N' -tris(3-methyl-pyridin-2-ylmethyl)-ethylenediamine;
 N -trimethylammoniumpropyl- N,N',N' -tris(pyridin-2-ylmethyl)-ethylenediamine;
 N -(2-hydroxyethylene)- N,N',N' -tris(pyridin-2-ylmethyl)-ethylenediamine;
 N,N,N',N' -tetrakis(3-methyl-pyridin-2-ylmethyl)-ethylenediamine;
 N,N' -dimethyl- N,N' -bis(pyridin-2-ylmethyl)-cyclohexane-1,2-diamine;
 N -(2-hydroxyethylene)- N,N',N' -tris(3-methyl-pyridin-2-ylmethyl)-ethylenediamine;

30

- N-methyl-N,N',N'-tris(pyridin-2-ylmethyl)-ethylenediamine;
N-methyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)-
ethylenediamine;
N-methyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)-
5 ethylenediamine;
N-methyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)-
ethylenediamine;
N-benzyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)-
ethylenediamine;
10 N-ethyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)-
ethylenediamine;
N,N,N'-tris(3-methyl-pyridin-2-ylmethyl)-N'(2'-methoxy-
ethyl-1)-ethylenediamine;
N,N,N'-tris(1-methyl-benzimidazol-2-yl)-N'-methyl-
15 ethylenediamine;
N-(furan-2-yl)-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)-
ethylenediamine;
N-(2-hydroxyethylene)-N,N',N'-tris(3-ethyl-pyridin-2-
ylmethyl)-ethylenediamine;
20 N-methyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;
N-ethyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;
25 N-benzyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;
N-(2-hydroxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine;
N-(2-methoxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-
30 ylmethyl)ethylene-1,2-diamine;

N-methyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-ethyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

5 N-benzyl-N,N',N'-tris(5-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-(2-hydroxyethyl)-N,N',N'-tris(5-methyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine;

10 N-(2-methoxyethyl)-N,N',N'-tris(5-methyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine;

N-methyl-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

15 N-ethyl-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-benzyl-N,N',N'-tris(3-ethyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-(2-hydroxyethyl)-N,N',N'-tris(3-ethyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine;

20 N-(2-methoxyethyl)-N,N',N'-tris(3-ethyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine;

N-methyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

25 N-ethyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-benzyl-N,N',N'-tris(5-ethyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine; and

30 N-(2-methoxyethyl)-N,N',N'-tris(5-ethyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine.

More preferred ligands are:

N-methyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;


5 N-ethyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-benzyl-N,N',N'-tris(3-methyl-pyridin-2-ylmethyl)ethylene-
1,2-diamine;

N-(2-hydroxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine; and

10 N-(2-methoxyethyl)-N,N',N'-tris(3-methyl-pyridin-2-
ylmethyl)ethylene-1,2-diamine.

(C) Ligands of the general formula (IC):




(IC)

wherein

20 Z_1 , Z_2 and Z_3 independently represent a coordinating
group selected from carboxylate, amido, $-NH-C(NH)NH_2$,
hydroxyphenyl, an optionally substituted heterocyclic ring
or an optionally substituted heteroaromatic ring selected
from pyridine, pyrimidine, pyrazine, pyrazole, imidazole,
benzimidazole, quinoline, quinoxaline, triazole,
isoquinoline, carbazole, indole, isoindole, oxazole and
25 thiazole;

Q₁, Q₂, and Q₃ independently represent a group of the formula:



5

wherein

$5 \geq a+b+c \geq 1$; $a=0-5$; $b=0-5$; $c=0-5$; $n=1$ or 2 ;

10

Y independently represents a group selected from -O-, -S-, -SO-, -SO₂-, -C(O)-, arylene, alkylene, heteroarylene, heterocycloalkylene, -(G)P-, -P(O)- and -(G)N-, wherein G is selected from hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, each except hydrogen being optionally

15

substituted by one or more functional groups E; and

20

R5, R6, R7, R8 independently represent a group selected from hydrogen, hydroxyl, halogen, -R and -OR, wherein R represents alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or a carbonyl derivative group, R being optionally substituted by one or more functional groups E, or R5 together with R6, or R7 together with R8, or both, represent oxygen,

25

or R5 together with R7 and/or independently R6 together with R8, or R5 together with R8 and/or independently R6

together with R7, represent C₁₋₆-alkylene optionally substituted by C₁₋₄-alkyl, -F, -Cl, -Br or -I.

5 Z₁, Z₂ and Z₃ each represent a coordinating group, preferably selected from optionally substituted pyridin-2-yl, optionally substituted imidazol-2-yl, optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl. Preferably, Z₁, Z₂ and Z₃ each represent optionally substituted pyridin-2-yl.

10

Optional substituents for the groups Z₁, Z₂ and Z₃ are preferably selected from C₁₋₄-alkyl, aryl, arylalkyl, heteroaryl, methoxy, hydroxy, nitro, amino, carboxyl, halo, and carbonyl, preferably methyl.

15

Also preferred is that Q₁, Q₂ and Q₃ are defined such that a=b=0, c=1 or 2, and n=1.

Preferably, each Q₁, Q₂ and Q₃ independently represent C₁₋₄-alkylene, more preferably a group selected from -CH₂- and -CH₂CH₂-.

20


The groups R5, R6, R7, R8 preferably independently represent a group selected from -H, hydroxy-C₀-C₂₀-alkyl, halo-C₀-C₂₀-alkyl, nitroso, formyl-C₀-C₂₀-alkyl, carboxyl-C₀-C₂₀-alkyl and esters and salts thereof, carbamoyl-C₀-C₂₀-alkyl, sulfo-C₀-C₂₀-alkyl and esters and salts thereof, sulfamoyl-C₀-C₂₀-alkyl, amino-C₀-C₂₀-alkyl, aryl-C₀-C₂₀-alkyl, C₀-C₂₀-alkyl, alkoxy-C₀-C₈-alkyl, carbonyl-C₀-C₆-alkoxy, and C₀-C₂₀-alkylamide. Preferably, none of R5-R8 is linked together.

30

Preferably, the ligand is selected from tris(pyridin-2-ylmethyl)amine, tris(3-methyl-pyridin-2-ylmethyl)amine, tris(5-methyl-pyridin-2-ylmethyl)amine, and tris(6-methyl-pyridin-2-ylmethyl)amine.

5

(D) Ligands of the general formula (ID):



10

(ID)

wherein


R_1 , R_2 , and R_3 independently represent a group selected from hydrogen, hydroxyl, halogen, $-NH-C(NH)NH_2$, $-R$ and $-OR$, wherein R = alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or a carbonyl derivative group, R being optionally substituted by one or more functional groups E ;

20

Q independently represent a group selected from C_{2-3} -alkylene optionally substituted by H , benzyl or C_{1-8} -alkyl;

Q_1 , Q_2 and Q_3 independently represent a group of the formula:

25



wherein

5 $5 \geq a+b+c \geq 1$; $a=0-5$; $b=0-5$; $c=0-5$; $n=1$ or 2 ;

Y independently represents a group selected from -O-, -S-, -SO-, -SO₂-, -C(O)-, arylene, alkylene, heteroarylene, heterocycloalkylene, -(G)P-, -P(O)- and -(G)N-, wherein G
10 is selected from hydrogen, alkyl, aryl, arylalkyl, cycloalkyl, each except hydrogen being optionally substituted by one or more functional groups E; and

R5, R6, R7, R8 independently represent a group selected
15 from hydrogen, hydroxyl, halogen, -R and -OR, wherein R represents alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or a carbonyl derivative group, R being optionally substituted by one or more functional groups E,
or R5 together with R6, or R7 together with R8, or
20 both, represent oxygen,

or R5 together with R7 and/or independently R6 together
with R8, or R5 together with R8 and/or independently R6
together with R7, represent C₁₋₆-alkylene optionally
substituted by C₁₋₄-alkyl, -F, -Cl, -Br or -I,

provided that at least one, preferably at least two, of R_1 , R_2 and R_3 is a coordinating group.

At least two, and preferably at least three, of R_1 , R_2 and R_3 independently represent a group selected from carboxylate, amido, $-\text{NH}-\text{C}(\text{NH})\text{NH}_2$, hydroxyphenyl, an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole. Preferably, at least two of R_1 , R_2 , R_3 each independently represent a coordinating group selected from optionally substituted pyridin-2-yl, optionally substituted imidazol-2-yl, optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl.


Preferably, substituents for groups R_1 , R_2 , R_3 , when representing a heterocyclic or heteroaromatic ring, are selected from C_{1-4} -alkyl, aryl, arylalkyl, heteroaryl, methoxy, hydroxy, nitro, amino, carboxyl, halo, and carbonyl.

Preferably, Q_1 , Q_2 and Q_3 are defined such that $a=b=0$, $c=1,2,3$ or 4 and $n=1$. Preferably, the groups Q_1 , Q_2 and Q_3 independently represent a group selected from $-\text{CH}_2-$ and $-\text{CH}_2\text{CH}_2-$.

Group Q is preferably a group selected from $-\text{CH}_2\text{CH}_2-$ and $-\text{CH}_2\text{CH}_2\text{CH}_2-$.

The groups R5, R6, R7, R8 preferably independently represent a group selected from -H, hydroxy-C₀-C₂₀-alkyl, halo-C₀-C₂₀-alkyl, nitroso, formyl-C₀-C₂₀-alkyl, carboxyl-C₀-C₂₀-alkyl and esters and salts thereof, carbamoyl-C₀-C₂₀-alkyl, sulfo-C₀-C₂₀-alkyl and esters and salts thereof, sulfamoyl-C₀-C₂₀-alkyl, amino-C₀-C₂₀-alkyl, aryl-C₀-C₂₀-alkyl, C₀-C₂₀-alkyl, alkoxy-C₀-C₈-alkyl, carbonyl-C₀-C₆-alkoxy, and C₀-C₂₀-alkylamide. Preferably, none of R5-R8 is linked together.

10 In a preferred aspect, the ligand is of the general formula (IID):



(IID)

15 wherein R1, R2, R3 are as defined previously for R₁, R₂, R₃, and Q₁, Q₂, Q₃ are as defined previously.

Preferred classes of ligands according to this preferred aspect, as represented by formula (IID) above, are as follows:

(i) ligands of the general formula (IID) wherein:

20 R1, R2, R3 each independently represent a coordinating group selected from carboxylate, amido, -NH-C(NH)NH₂,
25

hydroxyphenyl, an optionally substituted heterocyclic ring or an optionally substituted heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole,
5 isoquinoline, carbazole, indole, isoindole, oxazole and thiazole.

In this class, we prefer that:

R1, R2, R3 each independently represent a coordinating
10 group selected from optionally substituted pyridin-2-yl, optionally substituted imidazol-2-yl, optionally substituted imidazol-4-yl, optionally substituted pyrazol-1-yl, and optionally substituted quinolin-2-yl.

15 (ii) ligands of the general formula (IID) wherein:

two of R1, R2, R3 each independently represent a coordinating group selected from carboxylate, amido, -NH-C(NH)NH₂, hydroxyphenyl, an optionally substituted heterocyclic ring or an optionally substituted
20 heteroaromatic ring selected from pyridine, pyrimidine, pyrazine, pyrazole, imidazole, benzimidazole, quinoline, quinoxaline, triazole, isoquinoline, carbazole, indole, isoindole, oxazole and thiazole; and

one of R1, R2, R3 represents a group selected from
25 hydrogen, C₁₋₂₀ optionally substituted alkyl, C₁₋₂₀ optionally substituted arylalkyl, aryl, and C₁₋₂₀ optionally substituted NR₃⁺ (wherein R=C₁₋₈-alkyl).


In this class, we prefer that:

30 two of R1, R2, R3 each independently represent a coordinating group selected from optionally substituted

pyridin-2-yl, optionally substituted imidazol-2-yl,
optionally substituted imidazol-4-yl, optionally substituted
pyrazol-1-yl, and optionally substituted quinolin-2-yl; and


one of R1, R2, R3 represents a group selected from
5 hydrogen, C₁₋₁₀ optionally substituted alkyl, C₁₋₅-furanyl, C₁₋₅
optionally substituted benzylalkyl, benzyl, C₁₋₅ optionally
substituted alkoxy, and C₁₋₂₀ optionally substituted N⁺Me₃.

10 In especially preferred embodiments, the ligand is selected
from:



wherein -Et represents ethyl, -Py represents pyridin-2-yl,
15 Pz3 represents pyrazol-3-yl, Pz1 represents pyrazol-1-yl,
and Qu represents quinolin-2-yl.

(E) Ligands of the general formula (IE):



(IE)


wherein

g represents zero or an integer from 1 to 6;

10 r represents an integer from 1 to 6;

s represents zero or an integer from 1 to 6;

Q1 and Q2 independently represent a group of the formula:



wherein

20 $5 \geq d+e+f \geq 1$; $d=0-5$; $e=0-5$; $f=0-5$;

each Y1 independently represents a group selected from -O-, -S-, -SO-, -SO₂-, -C(O)-, arylene, alkylene,

heteroarylene, heterocycloalkylene, -(G)P-, -P(O)- and -

(G)N-, wherein G is selected from hydrogen, alkyl, aryl,

25 arylalkyl, cycloalkyl, each except hydrogen being optionally substituted by one or more functional groups E;

if $s > 1$, each $-\text{N}(\text{R1})-(\text{Q1})_r-$ group is independently defined;

R1, R2, R6, R7, R8, R9 independently represent a group selected from hydrogen, hydroxyl, halogen, -R and -OR, wherein R represents alkyl, alkenyl, cycloalkyl, heterocycloalkyl, aryl, heteroaryl or a carbonyl derivative group, R being optionally substituted by one or more functional groups E,

or R6 together with R7, or R8 together with R9, or both, represent oxygen,

or R6 together with R8 and/or independently R7 together with R9, or R6 together with R9 and/or independently R7 together with R8, represent C₁₋₆-alkylene optionally substituted by C₁₋₄-alkyl, -F, -Cl, -Br or -I;

or one of R1-R9 is a bridging group bound to another moiety of the same general formula;

T1 and T2 independently represent groups R4 and R5, wherein R4 and R5 are as defined for R1-R9, and if g=0 and s>0, R1 together with R4, and/or R2 together with R5, may optionally independently represent =CH-R10, wherein R10 is as defined for R1-R9, or

T1 and T2 may together (-T2-T1-) represent a covalent bond linkage when s>1 and g>0;

if T1 and T2 together represent a single bond linkage, Q1 and/or Q2 may independently represent a group of the formula: =CH-[-Y1-]_e-CH= provided R1 and/or R2 are absent, and R1 and/or R2 may be absent provided Q1 and/or Q2 independently represent a group of the formula: =CH-[-Y1-]_e-CH=.

The groups R1-R9 are preferably independently selected from -H, hydroxy-C₀-C₂₀-alkyl, halo-C₀-C₂₀-alkyl, nitroso, formyl-C₀-C₂₀-alkyl, carboxyl-C₀-C₂₀-alkyl and esters and salts thereof, carbamoyl-C₀-C₂₀-alkyl, sulpho-C₀-C₂₀-alkyl and


5 esters and salts thereof, sulphamoyl-C₀-C₂₀-alkyl, amino-C₀-C₂₀-alkyl, aryl-C₀-C₂₀-alkyl, heteroaryl-C₀-C₂₀-alkyl, C₀-C₂₀-alkyl, alkoxy-C₀-C₈-alkyl, carbonyl-C₀-C₆-alkoxy, and aryl-C₀-C₆-alkyl and C₀-C₂₀-alkylamide.

10 One of R1-R9 may be a bridging group which links the ligand moiety to a second ligand moiety of preferably the same general structure. In this case the bridging group is independently defined according to the formula for Q1, Q2, preferably being alkylene or hydroxy-alkylene or a

15 heteroaryl-containing bridge, more preferably C₁₋₆-alkylene optionally substituted by C₁₋₄-alkyl, -F, -Cl, -Br or -I.

In a first variant according to formula (IE), the groups T1 and T2 together form a single bond linkage and s>1,

20 according to general formula (IIE):




(IIE)

wherein R3 independently represents a group as defined for R1-R9; Q3 independently represents a group as defined for Q1, Q2; h represents zero or an integer from 1 to 6; and $\underline{s}=s-1$.

5

In a first embodiment of the first variant, in general formula (IIE), $\underline{s}=1, 2$ or 3 ; $r=g=h=1$; $d=2$ or 3 ; $e=f=0$; $R6=R7=H$, preferably such that the ligand has a general formula selected from:


11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197



- 20 In these preferred examples, R1, R2, R3 and R4 are preferably independently selected from -H, alkyl, aryl, heteroaryl, and/or one of R1-R4 represents a bridging group bound to another moiety of the same general formula and/or
- 25 linking N atoms in the same moiety, with the bridging group being alkylene or hydroxy-alkylene or a heteroaryl-containing bridge, preferably heteroarylene. More preferably, R1, R2, R3 and R4 are independently selected from -H, methyl, ethyl, isopropyl, nitrogen-containing
- 30 heteroaryl, or a bridging group bound to another moiety of the same general formula or linking N atoms in the same


moiety with the bridging group being alkylene or hydroxy-alkylene.

In a second embodiment of the first variant, in general
 5 formula (IIE), $s=2$ and $r=g=h=1$, according to the general formula:



In this second embodiment, preferably R1-R4 are absent; both
 10 Q1 and Q3 represent $=CH-[-Y1-]_e-CH=$; and both Q2 and Q4 represent $-CH_2-[-Y1-]_n-CH_2-$.

Thus, preferably the ligand has the general formula:




15

wherein A represents optionally substituted alkylene optionally interrupted by a heteroatom; and n is zero or an integer from 1 to 5.

Preferably, R1-R6 represent hydrogen, n=1 and A= -CH₂-, -CHOH-, -CH₂N(R)CH₂- or -CH₂CH₂N(R)CH₂CH₂- wherein R represents hydrogen or alkyl, more preferably A= -CH₂-, -CHOH- or -CH₂CH₂NHCH₂CH₂-.


5

In a second variant according to formula (IE), T1 and T2 independently represent groups R4, R5 as defined for R1-R9, according to the general formula (IIIE):



In a first embodiment of the second variant, in general formula (IIIE), s=1; r=1; g=0; d=f=1; e=0-4; Y1= -CH₂-; and R1 together with R4, and/or R2 together with R5, independently represent =CH-R10, wherein R10 is as defined for R1-R9. In one example, R2 together with R5 represents =CH-R10, with R1 and R4 being two separate groups.

Alternatively, both R1 together with R4, and R2 together with R5 may independently represent =CH-R10. Thus, preferred ligands may for example have a structure selected from:



25


wherein n = 0-4.

Preferably, the ligand is selected from:



- 5 wherein R1 and R2 are selected from optionally substituted phenols, heteroaryl-C₀-C₂₀-alkyls, R3 and R4 are selected from -H, alkyl, aryl, optionally substituted phenols, heteroaryl-C₀-C₂₀-alkyls, alkylaryl, aminoalkyl, alkoxy, more preferably R1 and R2 being selected from optionally substituted phenols, heteroaryl-C₀-C₂-alkyls, R3 and R4 are selected from -H, alkyl, aryl, optionally substituted phenols, nitrogen-heteroaryl-C₀-C₂-alkyls.


15 In a second embodiment of the second variant, in general formula (IIIE), s=1; r=1; g=0; d=f=1; e=1-4; Y1= -C(R')(R''), wherein R' and R'' are independently as defined for R1-R9. Preferably, the ligand has the general formula:



- 20 The groups R1, R2, R3, R4, R5 in this formula are preferably -H or C₀-C₂₀-alkyl, n=0 or 1, R6 is -H, alkyl, -OH or -SH, and R7, R8, R9, R10 are preferably each independently selected from -H, C₀-C₂₀-alkyl, heteroaryl-C₀-C₂₀-alkyl, alkoxy-C₀-C₈-alkyl and amino-C₀-C₂₀-alkyl.

In a third embodiment of the second variant, in general formula (IIIE), $s=0$; $g=1$; $d=e=0$; $f=1-4$. Preferably, the ligand has the general formula:


5



10

This class of ligand is particularly preferred according to the invention.


More preferably, the ligand has the general formula:



15

wherein R^1 , R^2 , R^3 are as defined for R^2 , R^4 , R^5 .

In a fourth embodiment of the second variant, the ligand is a pentadentate ligand of the general formula (IVE):



20

(IVE)

wherein

each R^1 , R^2 independently represents $-\text{R}^4-\text{R}^5$,

R^3 represents hydrogen, optionally substituted alkyl, aryl or arylalkyl, or $-R^4-R^5$,

each R^4 independently represents a single bond or optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene, alkylene ether, carboxylic ester or carboxylic amide, and

each R^5 independently represents an optionally N-substituted aminoalkyl group or an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl.

Ligands of the class represented by general formula (IVE) are also particularly preferred according to the invention.

The ligand having the general formula (IVE), as defined above, is a pentadentate ligand. By 'pentadentate' herein is meant that five hetero atoms can coordinate to the metal M ion in the metal-complex.

In formula (IVE), one coordinating hetero atom is provided by the nitrogen atom in the methylamine backbone, and preferably one coordinating hetero atom is contained in each of the four R^1 and R^2 side groups. Preferably, all the coordinating hetero atoms are nitrogen atoms.

The ligand of formula (IVE) preferably comprises at least two substituted or unsubstituted heteroaryl groups in the four side groups. The heteroaryl group is preferably a pyridin-2-yl group and, if substituted, preferably a methyl- or ethyl-substituted pyridin-2-yl group. More preferably, the heteroaryl group is an unsubstituted pyridin-2-yl group.

Preferably, the heteroaryl group is linked to methylamine, and preferably to the N atom thereof, via a methylene group. Preferably, the ligand of formula (IVE) contains at least one optionally substituted amino-alkyl side group, more preferably two amino-ethyl side groups, in particular 2-(N-alkyl)amino-ethyl or 2-(N,N-dialkyl)amino-ethyl.

Thus, in formula (IVE) preferably R^1 represents pyridin-2-yl or R^2 represents pyridin-2-yl-methyl. Preferably R^2 or R^1 represents 2-amino-ethyl, 2-(N-(m)ethyl)amino-ethyl or 2-(N,N-di(m)ethyl)amino-ethyl. If substituted, R^5 preferably represents 3-methyl pyridin-2-yl. R^3 preferably represents hydrogen, benzyl or methyl.

Examples of preferred ligands of formula (IVE) in their simplest forms are:

(i) pyridin-2-yl containing ligands such as:

N,N-bis(pyridin-2-yl-methyl)-bis(pyridin-2-yl)methylamine;
N,N-bis(pyrazol-1-yl-methyl)-bis(pyridin-2-yl)methylamine;
N,N-bis(imidazol-2-yl-methyl)-bis(pyridin-2-yl)methylamine;
N,N-bis(1,2,4-triazol-1-yl-methyl)-bis(pyridin-2-yl)methylamine;
N,N-bis(pyridin-2-yl-methyl)-bis(pyrazol-1-yl)methylamine;
N,N-bis(pyridin-2-yl-methyl)-bis(imidazol-2-yl)methylamine;
N,N-bis(pyridin-2-yl-methyl)-bis(1,2,4-triazol-1-yl)methylamine;
N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;
N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyrazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(pyrazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

5 N,N-bis(imidazol-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(imidazol-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

10 N,N-bis(1,2,4-triazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(1,2,4-triazol-1-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyrazol-1-yl)-1-aminoethane;

15 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyrazol-1-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(imidazol-2-yl)-1-aminoethane;

20 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(imidazol-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(1,2,4-triazol-1-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(1,2,4-triazol-1-yl)-1-aminoethane;

25 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane;

30 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(4-sulphonic acid-phenyl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(pyridin-2-yl)-1-aminoethane;

5 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(pyridin-3-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(pyridin-4-yl)-1-aminoethane;

10 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(1-alkyl-pyridinium-4-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(1-alkyl-pyridinium-3-yl)-1-aminoethane;

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-(1-alkyl-pyridinium-2-yl)-1-aminoethane;

15

(ii) 2-amino-ethyl containing ligands such as:

N,N-bis(2-(N-alkyl)amino-ethyl)-bis(pyridin-2-yl)methylamine;

20 N,N-bis(2-(N-alkyl)amino-ethyl)-bis(pyrazol-1-yl)methylamine;

N,N-bis(2-(N-alkyl)amino-ethyl)-bis(imidazol-2-yl)methylamine;

N,N-bis(2-(N-alkyl)amino-ethyl)-bis(1,2,4-triazol-1-yl)methylamine;

25 N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(pyridin-2-yl)methylamine;

N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(pyrazol-1-yl)methylamine;

30 N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(imidazol-2-yl)methylamine;

N,N-bis(2-(N,N-dialkyl)amino-ethyl)-bis(1,2,4-triazol-1-yl)methylamine;

N,N-bis(pyridin-2-yl-methyl)-bis(2-amino-ethyl)methylamine;

N,N-bis(pyrazol-1-yl-methyl)-bis(2-amino-ethyl)methylamine;

5 N,N-bis(imidazol-2-yl-methyl)-bis(2-amino-ethyl)methylamine;

N,N-bis(1,2,4-triazol-1-yl-methyl)-bis(2-amino-ethyl)methylamine.

More preferred ligands are:


10 N,N-bis(pyridin-2-yl-methyl)-bis(pyridin-2-yl)methylamine, hereafter referred to as N4Py.

N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, hereafter referred to as MeN4Py,

15 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-2-phenyl-1-aminoethane, hereafter referred to as BzN4Py.

In a fifth embodiment of the second variant, the ligand represents a pentadentate or hexadentate ligand of general formula (VE):

20



(VE)

wherein

25 each R^1 independently represents $-R^3-V$, in which R^3 represents optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene or alkylene ether, and V represents an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, 30 imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl;

W represents an optionally substituted alkylene bridging group selected from -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, -CH₂-C₆H₄-CH₂-, -CH₂-C₆H₁₀-CH₂-, and -CH₂-C₁₀H₆-CH₂-; and

5 R² represents a group selected from R¹, and alkyl, aryl and arylalkyl groups optionally substituted with a substituent selected from hydroxy, alkoxy, phenoxy, carboxylate, carboxamide, carboxylic ester, sulphonate, amine, alkylamine and N⁺(R⁴)₃, wherein R⁴ is selected from
10 hydrogen, alkanyl, alkenyl, arylalkanyl, arylalkenyl, oxyalkanyl, oxyalkenyl, aminoalkanyl, aminoalkenyl, alkanyl ether and alkenyl ether.

The ligand having the general formula (VE), as defined
15 above, is a pentadentate ligand or, if R¹=R², can be a hexadentate ligand. As mentioned above, by 'pentadentate' is meant that five hetero atoms can coordinate to the metal M ion in the metal-complex. Similarly, by 'hexadentate' is meant that six hetero atoms can in principle coordinate to
20 the metal M ion. However, in this case it is believed that one of the arms will not be bound in the complex, so that the hexadentate ligand will be penta coordinating.

In the formula (VE), two hetero atoms are linked by the
25 bridging group W and one coordinating hetero atom is contained in each of the three R¹ groups. Preferably, the coordinating hetero atoms are nitrogen atoms.

The ligand of formula (VE) comprises at least one optionally
30 substituted heteroaryl group in each of the three R¹ groups. Preferably, the heteroaryl group is a pyridin-2-yl group, in

particular a methyl- or ethyl-substituted pyridin-2-yl group. The heteroaryl group is linked to an N atom in formula (VE), preferably via an alkylene group, more preferably a methylene group. Most preferably, the
5 heteroaryl group is a 3-methyl-pyridin-2-yl group linked to an N atom via methylene.

The group R^2 in formula (VE) is a substituted or unsubstituted alkyl, aryl or arylalkyl group, or a group R^1 .
10 However, preferably R^2 is different from each of the groups R^1 in the formula above. Preferably, R^2 is methyl, ethyl, benzyl, 2-hydroxyethyl or 2-methoxyethyl. More preferably, R^2 is methyl or ethyl.

15 The bridging group W may be a substituted or unsubstituted alkylene group selected from $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2-\text{C}_6\text{H}_4-\text{CH}_2-$, $-\text{CH}_2-\text{C}_6\text{H}_{10}-\text{CH}_2-$, and $-\text{CH}_2-\text{C}_{10}\text{H}_6-\text{CH}_2-$ (wherein $-\text{C}_6\text{H}_4-$, $-\text{C}_6\text{H}_{10}-$, $-\text{C}_{10}\text{H}_6-$ can be *ortho*-, *para*-, or *meta*- C_6H_4- , $-\text{C}_6\text{H}_{10}-$, $-\text{C}_{10}\text{H}_6-$). Preferably, the bridging group
20 W is an ethylene or 1,4-butylene group, more preferably an ethylene group.

Preferably, V represents substituted pyridin-2-yl, especially methyl-substituted or ethyl-substituted pyridin-
25 2-yl, and most preferably V represents 3-methyl pyridin-2-yl.

(F) Ligands of the classes disclosed in WO-A-98/39098 and WO-A-98/39406.

The counter ions Y in formula (A1) balance the charge z on the complex formed by the ligand L, metal M and coordinating species X. Thus, if the charge z is positive, Y may be an anion such as RCOO^- , BPh_4^- , ClO_4^- , BF_4^- , PF_6^- , RSO_3^- , RSO_4^- , SO_4^{2-} , NO_3^- , F^- , Cl^- , Br^- , or I^- , with R being hydrogen, optionally substituted alkyl or optionally substituted aryl. If z is negative, Y may be a common cation such as an alkali metal, alkaline earth metal or (alkyl)ammonium cation.

Suitable counter ions Y include those which give rise to the formation of storage-stable solids. Preferred counter ions for the preferred metal complexes are selected from R^7COO^- , ClO_4^- , BF_4^- , PF_6^- , RSO_3^- (in particular CF_3SO_3^-), RSO_4^- , SO_4^{2-} , NO_3^- , F^- , Cl^- , Br^- , and I^- , wherein R represents hydrogen or optionally substituted phenyl, naphthyl or C_1 - C_4 alkyl.

It will be appreciated that the complex (A1) can be formed by any appropriate means, including *in situ* formation whereby precursors of the complex are transformed into the active complex of general formula (A1) under conditions of storage or use. Preferably, the complex is formed as a well-defined complex or in a solvent mixture comprising a salt of the metal M and the ligand L or ligand L-generating species. Alternatively, the catalyst may be formed *in situ* from suitable precursors for the complex, for example in a solution or dispersion containing the precursor materials. In one such example, the active catalyst may be formed *in situ* in a mixture comprising a salt of the metal M and the ligand L, or a ligand L-generating species, in a suitable solvent. Thus, for example, if M is iron, an iron salt such as FeSO_4 can be mixed in solution with the ligand L, or a

ligand L-generating species, to form the active complex.

Thus, for example, the composition may be formed from a mixture of the ligand L and a metal salt MX_n in which preferably $n=1-5$, more preferably 1-3. In another such example, the ligand

5 L, or a ligand L-generating species, can be mixed with metal M ions present in the substrate or wash liquor to form the active catalyst *in situ*. Suitable ligand L-generating species include metal-free compounds or metal coordination complexes that comprise the ligand L and can be substituted
10 by metal M ions to form the active complex according to the formula (A1).

Throughout the description and claims generic groups have been used, for example alkyl, alkoxy, aryl. Unless

15 otherwise specified the following are preferred group restrictions that may be applied to generic groups found within compounds disclosed herein:

20 alkyl: C1-C6-alkyl,

alkenyl: C2-C6-alkenyl,

cycloalkyl: C3-C8-cycloalkyl,

25 alkoxy: C1-C6-alkoxy,

alkylene: selected from the group consisting of: methylene;
1,1-ethylene; 1,2-ethylene; 1,1-propylene; 1,2-propylene;
1,3-propylene; 2,2-propylene; butan-2-ol-1,4-diyl; propan-2-
30 ol-1,3-diyl; and 1,4-butylene,

aryl: selected from homoaromatic compounds having a molecular weight under 300,

arylene: selected from the group consisting of: 1,2-benzene; 1,3-benzene; 1,4-benzene; 1,2-naphthalene; 1,3-naphthalene; 1,4-naphthalene; 2,3-naphthalene; phenol-2,3-diyl; phenol-2,4-diyl; phenol-2,5-diyl; and phenol-2,6-diyl,

10 heteroaryl: selected from the group consisting of: pyridinyl; pyrimidinyl; pyrazinyl; triazolyl, pyridazinyl; 1,3,5-triazinyl; quinolinyl; isoquinolinyl; quinoxalinyl; imidazolyl; pyrazolyl; benzimidazolyl; thiazolyl; oxazolidinyl; pyrrolyl; carbazolyl; indolyl; and isoindolyl,

15 heteroarylene: selected from the group consisting of: pyridin-2,3-diyl; pyridin-2,4-diyl; pyridin-2,5-diyl; pyridin-2,6-diyl; pyridin-3,4-diyl; pyridin-3,5-diyl; quinolin-2,3-diyl; quinolin-2,4-diyl; quinolin-2,8-diyl; 20 isoquinolin-1,3-diyl; isoquinolin-1,4-diyl; pyrazol-1,3-diyl; pyrazol-3,5-diyl; triazole-3,5-diyl; triazole-1,3-diyl; pyrazin-2,5-diyl; and imidazole-2,4-diyl,

heterocycloalkyl: selected from the group consisting of: pyrrolinyl; pyrrolidinyl; morpholinyl; piperidinyl; 25 piperazinyl; hexamethylene imine; and oxazolidinyl,

amine: the group $-N(R)_2$ wherein each R is independently selected from: hydrogen; C1-C6-alkyl; C1-C6-alkyl-C6H5; and phenyl, wherein when both R are C1-C6-alkyl both R together 30 may form an -NC3 to an -NC5 heterocyclic ring with any

remaining alkyl chain forming an alkyl substituent to the heterocyclic ring,

5 halogen: selected from the group consisting of: F; Cl; Br and I,

sulphonate: the group $-S(O)_2OR$, wherein R is selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C₆H₅; Li; Na; K; Cs; Mg; and Ca,

10

sulphate: the group $-OS(O)_2OR$, wherein R is selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C₆H₅; Li; Na; K; Cs; Mg; and Ca,

15 sulphone: the group $-S(O)_2R$, wherein R is selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C₆H₅ and amine (to give sulphonamide) selected from the group: $-NR'_2$, wherein each R' is independently selected from: hydrogen; C1-C6-alkyl; C1-C6-alkyl-C₆H₅; and phenyl, wherein when both
20 R' are C1-C6-alkyl both R' together may form an -NC₃ to an -NC₅ heterocyclic ring with any remaining alkyl chain forming an alkyl substituent to the heterocyclic ring,

carboxylate derivative: the group $-C(O)OR$, wherein R is
25 selected from: hydrogen, C1-C6-alkyl; phenyl; C1-C6-alkyl-C₆H₅, Li; Na; K; Cs; Mg; and Ca,

carbonyl derivative: the group $-C(O)R$, wherein R is
30 selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C₆H₅ and amine (to give amide) selected from the group: $-NR'_2$, wherein each R' is independently selected from:

hydrogen; C1-C6-alkyl; C1-C6-alkyl-C6H5; and phenyl, wherein when both R' are C1-C6-alkyl both R' together may form an -NC3 to an -NC5 heterocyclic ring with any remaining alkyl chain forming an alkyl substituent to the heterocyclic ring,

5

phosphonate: the group $-P(O)(OR)_2$, wherein each R is independently selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C6H5; Li; Na; K; Cs; Mg; and Ca,

10 phosphate: the group $-OP(O)(OR)_2$, wherein each R is independently selected from: hydrogen; C1-C6-alkyl; phenyl; C1-C6-alkyl-C6H5; Li; Na; K; Cs; Mg; and Ca,

phosphine: the group $-P(R)_2$, wherein each R is
15 independently selected from: hydrogen; C1-C6-alkyl; phenyl; and C1-C6-alkyl-C6H5,

phosphine oxide: the group $-P(O)R_2$, wherein R is independently selected from: hydrogen; C1-C6-alkyl; phenyl; and C1-C6-alkyl-C6H5; and amine (to give phosphonamidate)
20 selected from the group: $-NR'_2$, wherein each R' is independently selected from: hydrogen; C1-C6-alkyl; C1-C6-alkyl-C6H5; and phenyl, wherein when both R' are C1-C6-alkyl both R' together may form an -NC3 to an -NC5 heterocyclic
25 ring with any remaining alkyl chain forming an alkyl substituent to the heterocyclic ring.

Unless otherwise specified the following are more preferred group restrictions that may be applied to groups found
30 within compounds disclosed herein:

alkyl: C1-C4-alkyl,

alkenyl: C3-C6-alkenyl,

5 cycloalkyl: C6-C8-cycloalkyl,

alkoxy: C1-C4-alkoxy,

alkylene: selected from the group consisting of: methylene;
10 1,2-ethylene; 1,3-propylene; butan-2-ol-1,4-diyl; and 1,4-butylene,

aryl: selected from group consisting of: phenyl;
biphenyl, naphthalenyl; anthracenyl; and phenanthrenyl,
15

arylene: selected from the group consisting of: 1,2-benzene, 1,3-benzene, 1,4-benzene, 1,2-naphthalene, 1,4-naphthalene, 2,3-naphthalene and phenol-2,6-diyl,

20 heteroaryl: selected from the group consisting of: pyridinyl; pyrimidinyl; quinolinyl; pyrazolyl; triazolyl; isoquinolinyl; imidazolyl; and oxazolidinyl,

heteroarylene: selected from the group consisting of:
25 pyridin-2,3-diyl; pyridin-2,4-diyl; pyridin-2,6-diyl; pyridin-3,5-diyl; quinolin-2,3-diyl; quinolin-2,4-diyl; isoquinolin-1,3-diyl; isoquinolin-1,4-diyl; pyrazol-3,5-diyl; and imidazole-2,4-diyl,

30 heterocycloalkyl: selected from the group consisting of: pyrrolidinyl; morpholinyl; piperidinyl; and piperazinyl,

amine: the group $-N(R)_2$, wherein each R is independently selected from: hydrogen; C1-C6-alkyl; and benzyl,

halogen: selected from the group consisting of: F and Cl,

5 sulphonate: the group $-S(O)_2OR$, wherein R is selected from: hydrogen; C1-C6-alkyl; Na; K; Mg; and Ca,

10 sulphate: the group $-OS(O)_2OR$, wherein R is selected from: hydrogen; C1-C6-alkyl; Na; K; Mg; and Ca,

15 sulphone: the group $-S(O)_2R$, wherein R is selected from: hydrogen; C1-C6-alkyl; benzyl and amine selected from the group: $-NR'_2$, wherein each R' is independently selected from: hydrogen; C1-C6-alkyl; and benzyl,

20 carboxylate derivative: the group $-C(O)OR$, wherein R is selected from hydrogen; Na; K; Mg; Ca; C1-C6-alkyl; and benzyl,

25 carbonyl derivative: the group: $-C(O)R$, wherein R is selected from: hydrogen; C1-C6-alkyl; benzyl and amine selected from the group: $-NR'_2$, wherein each R' is independently selected from: hydrogen; C1-C6-alkyl; and benzyl,

30 phosphonate: the group $-P(O)(OR)_2$, wherein each R is independently selected from: hydrogen; C1-C6-alkyl, benzyl; Na; K; Mg; and Ca,

phosphate: the group $-OP(O)(OR)_2$, wherein each R is independently selected from: hydrogen; C1-C6-alkyl; benzyl; Na; K; Mg; and Ca,

- 5 phosphine: the group $-P(R)_2$, wherein each R is independently selected from: hydrogen; C1-C6-alkyl; and benzyl,

- phosphine oxide: the group $-P(O)R_2$, wherein R is
10 independently selected from: hydrogen; C1-C6-alkyl; benzyl and amine selected from the group: $-NR'_2$, wherein each R' is independently selected from: hydrogen; C1-C6-alkyl; and benzyl.

- 15 Other compounds or ligands forming complexes with transition metals, and which are capable of catalysing bleaching by atmospheric oxygen, are suitable as organic substances in the liquid bleaching compositions of the present invention. These include the classes of complexes of a transition metal
20 coordinated to a macropolycyclic ligand disclosed in WO-A-98/39098 and WO-A-98/39406.

- The liquid bleaching compositions according to the present invention may be used for laundry cleaning, hard surface
25 cleaning (including cleaning of lavatories, kitchen work surfaces, floors, mechanical ware washing etc.). As is generally known in the art, bleaching compositions are also employed in waste-water treatment, pulp bleaching during the manufacture of paper, leather manufacture, dye transfer
30 inhibition, food processing, starch bleaching, sterilisation,

whitening in oral hygiene preparations and/or contact lens disinfection.

In the context of the present invention bleaching should be understood as relating generally to the decolourisation of stains or of other materials attached to or associated with a substrate. However, it is envisaged that the present invention can be applied where a requirement is the removal and/or neutralisation by an oxidative bleaching reaction of malodours or other undesirable components attached to or otherwise associated with a substrate. Furthermore, in the context of the present invention bleaching is to be understood as being restricted to any bleaching mechanism or process that does not require the presence of light or activation by light. Thus, photobleaching compositions and processes relying on the use of photobleach catalysts or photobleach activators and the presence of light are excluded from the present invention.

In typical washing compositions the level of the organic substance is such that the in-use level is from 0.05 μM to 50 mM, with preferred in-use levels for domestic laundry operations falling in the range 1 to 100 μM . Higher levels may be desired and applied in industrial bleaching processes, such as textile and paper pulp bleaching.

Preferably, the aqueous medium has a pH in the range from pH 6 to 13, more preferably from pH 6 to 11, and most preferably from 7 to 10.

The liquid bleaching composition of the present invention has particular application in detergent formulations,

especially for laundry cleaning. Accordingly, in another preferred embodiment, the present invention provides a liquid detergent bleach composition comprising a liquid bleaching composition as defined above and additionally a surface-active material, optionally together with detergency builder. In addition, the liquid bleaching composition may optionally contain soluble and non-soluble enzymes, enzyme stabiliser systems, functional polymers, polymers to modify the appearance and sensory properties of the liquid bleaching composition and optionally other minors such as a perfume or a fluorescer.

The liquid bleach composition according to the present invention may for example contain a surface-active material in an amount of from 10 to 50% by weight. The surface-active material may be naturally derived, such as soap, or a synthetic material selected from anionic, nonionic, amphoteric, zwitterionic, cationic actives and mixtures thereof. Many suitable actives are commercially available and are fully described in the literature, for example in "Surface Active Agents and Detergents", Volumes I and II, by Schwartz, Perry and Berch.

Typical synthetic anionic surface-actives are usually water-soluble alkali metal salts of organic sulphates and sulphonates having alkyl groups containing from about 8 to about 22 carbon atoms, the term "alkyl" being used to include the alkyl portion of higher aryl groups. Examples of suitable synthetic anionic detergent compounds are sodium and ammonium alkyl sulphates, especially those obtained by sulphating higher (C₈-C₁₈) alcohols produced, for example,

from tallow or coconut oil; sodium and ammonium alkyl (C₉-C₂₀) benzene sulphonates, particularly sodium linear secondary alkyl (C₁₀-C₁₅) benzene sulphonates; sodium alkyl glyceryl ether sulphates, especially those ethers of the

5 higher alcohols derived from tallow or coconut oil fatty acid monoglyceride sulphates and sulphonates; sodium and ammonium salts of sulphuric acid esters of higher (C₉-C₁₈) fatty alcohol alkylene oxide, particularly ethylene oxide, reaction products; the reaction products of fatty acids such

10 as coconut fatty acids esterified with isethionic acid and neutralised with sodium hydroxide; sodium and ammonium salts of fatty acid amides of methyl taurine; alkane monosulphonates such as those derived by reacting alpha-olefins (C₈-C₂₀) with sodium bisulphite and those derived by

15 reacting paraffins with SO₂ and Cl₂ and then hydrolysing with a base to produce a random sulphonate; sodium and ammonium (C₇-C₁₂) dialkyl sulphosuccinates; and olefin sulphonates, which term is used to describe material made by reacting olefins, particularly (C₁₀-C₂₀) alpha-olefins, with SO₃ and

20 then neutralising and hydrolysing the reaction product. The preferred anionic detergent compounds are sodium (C₁₀-C₁₅) alkylbenzene sulphonates, and sodium (C₁₆-C₁₈) alkyl ether sulphates.

25 Examples of suitable nonionic surface-active compounds which may be used, preferably together with the anionic surface-active compounds, include, in particular, the reaction products of alkylene oxides, usually ethylene oxide, with alkyl (C₆-C₂₂) phenols, generally 5-25 EO, i.e. 5-25 units of

30 ethylene oxides per molecule; and the condensation products of aliphatic (C₈-C₁₈) primary or secondary linear or branched

alcohols with ethylene oxide, generally 2-30 EO. Other so-called nonionic surface-actives include alkyl polyglycosides, sugar esters, long-chain tertiary amine oxides, long-chain tertiary phosphine oxides and dialkyl sulphoxides. The non-ionic surfactant liquid may be applied/ added in the form of a water-soluble sachet.

Amphoteric or zwitterionic surface-active compounds can also be used in the compositions of the invention. If any

10 amphoteric or zwitterionic detergent compounds are used, it is generally in small amounts in compositions based on the much more commonly used synthetic anionic and nonionic actives.

15 The liquid detergent bleach composition of the invention may comprise from 1 to 40 % wt of anionic surfactant and from 0 to 40 % by weight of nonionic surfactant. The liquid detergent may contain any mixture of non-ionic, anionic, cationic zwitterionic or combination thereof. Optionally, 20 fatty acid soaps (0-30%) may be present. The liquid detergent bleach composition of the present invention may also contains a detergency builder, for example in an amount of from about 5 to 80 % by weight, preferably from about 10 to 60 % by weight.

25

Builder materials may be selected from 1) calcium sequestrant materials, 2) precipitating materials, 3) calcium ion-exchange materials and 4) mixtures thereof.

Examples of calcium sequestrant builder materials include alkali metal polyphosphates, such as sodium tripolyphosphate; nitrilotriacetic acid and its water-soluble salts; the alkali metal salts of carboxymethyloxy succinic acid, ethylene diamine tetraacetic acid, oxydisuccinic acid, mellitic acid, benzene polycarboxylic acids, citric acid; and polyacetal carboxylates as disclosed in US-A-4,144,226 and US-A-4,146,495.

- 10 Examples of precipitating builder materials include sodium orthophosphate and sodium carbonate.

Examples of calcium ion-exchange builder materials include the various types of water-insoluble crystalline or amorphous aluminosilicates, of which zeolites are the best known representatives, e.g. zeolite A, zeolite B (also known as zeolite P), zeolite C, zeolite X, zeolite Y and also the zeolite P-type as described in EP-A-0,384,070.

- 20 In particular, the liquid bleaching compositions of the invention may contain any one of the organic and inorganic builder materials, though, for environmental reasons, phosphate builders are preferably omitted or only used in very small amounts. Typical builders usable in the present invention are, for example, sodium carbonate, calcite/carbonate, the sodium salt of nitrilotriacetic acid, sodium citrate, carboxymethyloxy malonate, carboxymethyloxy succinate and water-insoluble crystalline or amorphous aluminosilicate builder materials, each of which can be used as the main builder, either alone or in admixture with minor amounts of other builders or polymers as co-builder.

It is preferred that the liquid bleaching composition contains not more than 5% by weight of a carbonate builder, expressed as sodium carbonate, more preferably not more than 2.5 % by weight to substantially nil, if the composition pH
5 lies in the lower alkaline region of up to 10.

Apart from the components already mentioned, the liquid bleaching composition of the present invention can contain any of the conventional additives in amounts of which such
10 materials are normally employed in fabric washing detergent compositions. Examples of these additives include buffers such as carbonates, lather boosters, such as alkanolamides, particularly the monoethanol amides derived from palmkernel fatty acids and coconut fatty acids; lather depressants,
15 such as alkyl phosphates and silicones; anti-redeposition agents, such as sodium carboxymethyl cellulose and alkyl or substituted alkyl cellulose ethers; stabilisers, such as phosphonic acid derivatives (i.e. Dequest® types); fabric softening agents; inorganic salts and alkaline buffering
20 agents, such as sodium sulphate and sodium silicate; and, usually in very small amounts, fluorescent agents; perfumes; enzymes, such as proteases, cellulases, lipases, amylases and oxidases; germicides and colourants.

25 Transition metal sequestrants such as EDTA, and phosphonic acid derivatives such as EDTMP (ethylene diamine tetra(methylene phosphonate)) may also be included, in addition to the organic substance specified, for example to improve the stability sensitive ingredients such as enzymes,
30 fluorescent agents and perfumes, but provided the composition remains bleaching effective. However, the

liquid bleaching composition according to the present invention containing the organic substance, is preferably substantially, and more preferably completely, devoid of transition metal sequestrants (other than the organic substance).

Whilst the present invention is based on the catalytic bleaching of a substrate by atmospheric oxygen or air, it will be appreciated that small amounts of hydrogen peroxide or peroxy-based or -generating systems may be included in the liquid composition, if desired, provided that the chemical and physical stability of the composition is not thereby adversely affected to an unacceptable level. Therefore, by "substantially devoid of peroxygen bleach or peroxy-based or -generating bleach systems" is meant that the liquid bleaching composition contains from 0 to 50 %, preferably from 0 to 10 %, more preferably from 0 to 5 %, and optimally from 0 to 2 % by molar weight on an oxygen basis, of peroxygen bleach or peroxy-based or -generating bleach systems. Preferably, however, the liquid bleaching composition will be wholly devoid of peroxygen bleach or peroxy-based or -generating bleach systems.

Thus, at least 10 %, preferably at least 50 % and optimally at least 90 % of any bleaching of the substrate is effected by oxygen sourced from the air.

According to the fourth aspect, the organic substance in the liquid bleaching composition may be contacted to the textile fabric in any suitable manner. For example, it may be applied in a liquor that is then dried, for example as an

aqueous spray-on fabric treatment fluid or a wash liquor for laundry cleaning, or a non-aqueous dry cleaning fluid or spray-on aerosol fluid. Other suitable means of contacting the organic substance in liquid form to the textile may be
5 used, as further explained below.

Any suitable textile that is susceptible to bleaching or one that one might wish to subject to bleaching may be used. Preferably the textile is a laundry fabric or garment.

10

In a preferred embodiment of the fourth aspect, the method is carried out on a laundry fabric using an aqueous treatment liquor. In particular, the treatment may be effected in a wash cycle for cleaning laundry. More
15 preferably, the treatment is carried out in an aqueous detergent bleach wash liquid. In a preferred embodiment, the treated textile is dried, by allowing it to dry under ambient temperature or at elevated temperatures.

20 The bleaching method of the fourth aspect may be carried out by simply leaving the substrate in contact with the organic substance in the liquid bleaching composition for a sufficient period of time. Preferably, however, the organic substance is in an aqueous medium, and the aqueous medium on or containing
25 the substrate is agitated.

In a preferred embodiment of the fourth aspect, the treated textile is dried, by allowing it to dry under ambient temperature or at elevated temperatures.

30

In a particularly preferred embodiment the method according to the fourth aspect is carried out on a laundry fabric using aqueous treatment liquor. In particular the treatment may be effected in, or as an adjunct to, an essentially
5 conventional wash cycle for cleaning laundry. More preferably, the treatment is carried out in an aqueous detergent wash liquor. Preferably, the organic substance is delivered into the wash liquor from a liquid concentrate.

10 It is particularly advantageous that the organic substance in liquid composition used in the method of the fourth aspect makes use of atmospheric oxygen in its bleaching activity. This avoids the requirement that peroxygen bleaches and/or other relatively large quantities of
15 reactive substances need be used in the treatment process. Consequently, only a relatively small quantity of bleach active substance in liquid composition need be employed and this allows dosage routes to be exploited, which could previously not be used. Thus, while it is preferable to
20 include the organic substance in a liquid composition that is normally used in a washing process, such as a pre-treatment, main-wash, conditioning composition or ironing aid, other means for ensuring that the organic substance is present in the wash liquor may be envisaged.

25 For example, it is envisaged that the organic substance in the liquid composition can be presented in the form of a body from which it is slowly released during the whole or part of the laundry process. Such release can occur over the
30 course of a single wash or over the course of a plurality of washes. In the latter case it is envisaged that the organic

substance in liquid composition can be released from a carrier substrate used in association with the wash process, e.g. from a body placed in the dispenser drawer of a washing machine, elsewhere in the delivery system or in the drum of the washing machine. When used in the drum of the washing machine the carrier can be freely moving or fixed relative to the drum. Such fixing can be achieved by mechanical means, for example by barbs that interact with the drum wall, or employ other forces, for example a magnetic force.

The modification of a washing machine to provide for means to hold and retain such a carrier is envisaged similar means being known from the analogous art of toilet block manufacture. Freely moving carriers such as shuttles for dosage of surfactant materials and/or other detergent ingredients into the wash can comprise means for the release of the organic substance in the liquid composition into the wash.

In the alternative, the organic substance can be presented in the form of a liquid wash additive that preferably is soluble. Dosage of the additive can be unitary or in a quantity determined by the user. While it is envisaged that such additives can be used in the main washing cycle, the use of them in the conditioning or drying cycle is not hereby excluded.

The present invention is not limited to those circumstances in which a washing machine is employed, but can be applied where washing is performed in some alternative vessel. In these circumstances it is envisaged that the organic substance in liquid composition can be delivered by means of

slow release from the bowl, bucket or other vessel which is being employed, or from any implement which is being employed, such as a brush, bat or dolly, or from any suitable applicator for liquid compositions.

5

Suitable pre-treatment means for application of the organic substance from the liquid composition to the textile material prior to the main wash include sprays, pens, roller-ball devices and impregnated cloths or cloths :

10 containing microcapsules. Such means are well known in the analogous art of deodorant application and/or in spot treatment of textiles. Similar means for application are employed in those embodiments where the organic substance in liquid composition is applied after the main washing and/or
15 conditioning steps have been performed, e.g. prior to or after ironing or drying of the cloth. For example, the organic substance in liquid composition may be applied using tapes, sheets or sticking plasters coated or impregnated with the substance, or containing microcapsules of the
20 substance. The organic substance in liquid composition may for example be incorporated into a drier sheet so as to be activated or released during a tumble-drier cycle, or the organic substance in liquid composition can be provided in an impregnated or microcapsule-containing sheet so as to be
25 delivered to the textile when ironed.

The invention will now be further illustrated by way of the following non-limiting examples:

EXAMPLESExample 1

5 This example describes a synthesis of the catalyst as employed in Example 2:

(i) Preparation of MeN4Py ligand:

10 N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, MeN4Py, was prepared according to the procedure found in EP 0 909 809 A.

(ii) Synthesis of the complex FeMeN4PyCl₂:

15 MeN4Py ligand (33.7 g; 88.5mmoles) was dissolved in 500ml dry methanol. Small portions of FeCl₂.4H₂O (0.95eq; 16.7g; 84.0mmoles) were added, yielding a clear red solution. After addition, the solution was stirred for 30 minutes at room temperature, after which the methanol was removed (rotary-evaporator). The dry solid was ground and 150 ml of
20 ethylacetate was added and the mixture was stirred until a fine red powder was obtained. This powder was washed twice with ethyl acetate, dried in the air and further dried under vacuum (40 °C). El. Anal. Calc. for [Fe(MeN4py)Cl]Cl.2H₂O: C 53.03; H 5.16; N 12.89; Cl 13.07; Fe 10.01%. Found C 52.29/
25 52.03; H 5.05/5.03; N 12.55/12.61; Cl: 12.73/12.69; Fe: 10.06/10.01%.

Example 2:

30 Experiments with the FeMeN4PyCl₂ complex in a variety of liquid detergents were performed to establish bleaching

activity in various liquid detergent formulations and to determine stability upon storage.

FeMeN4PyCl₂ complex was added to several liquid detergent products and the stability and activity observed during storage.

The following commercially available liquid detergent compositions were used as base liquids: a) WISKTM liquid USA, 1999; b) OMOTM liquid NL, 1999; c) OMO-liquidoTM Brazil, 1999; and d) Rinse conditioner (RobijnTM - NL).

Incorporation of FeMeN4PyCl₂ in liquid detergents:

FeMeN4PyCl₂ was incorporated by post dosing a stock solution of 0.01 g/ml using an electrical stirrer (125 rpm, Heidolph RZR 2101). The final concentration in the product was 0.1% for all products. To the reference a same amount of water was added by post dosing to compensate for the post dose volume of the stock solution.

20

The activity of FeMeN4PyCl₂ was measured by washing tomato oil (TO) cloth samples in mini bottles for 15 minutes at a temperature of 25 °C and a dosage of 2 g/l product at 10 °FH. All of the liquids prepared were initially stable and homogeneous.

25

The following table lists compositions prepared. As detailed above base liquids a) to d) have had FeMeN4PyCl₂ incorporated therein. Compositions 5 to 8 are control liquids without added FeMeN4PyCl₂.

30

Composition No.	Liquids
1	Wisk™ liquid USA, 1999
2	OMO™ liquid NL, 1999
3	OMO-liquido™ Brazil, 1999
4	Rinse conditioner (Robijn™ - NL)
	Reference Liquids
5	Wisk™ liquid USA, 1999
6	OMO™ liquid NL, 1999
7	OMO™ -liquido™ Brazil, 1999
8	Rinse conditioner (Robijn™ - NL)

5 Cloth samples were washed in mini bottles with a liquid:cloth ratio of 1:20 and the samples were dried in a tumble dryer.

10 Bleaching activity was measured directly after the wash (after 2 hours), and after 1 one-day (24 hours) storage in the dark in order to establish post wash bleach effects. The five liquid formulations were stored under ambient conditions and the cleaning activity of the formulations without and with FeMeN4PyCl₂ was determined after certain periods of times. The times were immediately after preparation, and after 1, 2, 3, 4 and 6 weeks of storage. After the wash, the cloths were dried in a tumble drier and the reflectance was measured with a Minolta™ 3700d spectrophotometer at 460 nm. The difference in reflectance before and after the wash is defined as a ΔR_{460} value.

20 Tabulated results are shown in Tables 1 to 6 below.

Table 1

Directly after preparation	TO-stain ΔR 460		TO-stain ΔR 460	
10 FH, 2 g/l, T=25 C	2 hours after washing		1 day after washing	
Triplicate measurements	average	stdv	average	Stdv
Composition 5	14.3	1.6	26.6	3.1
Composition 1	16.5	1.4	34.6	0.6
Composition 6	12.9	0.7	20.0	2.9
Composition 2	17.2	1.4	35.7	0.8
Composition 7	16.2	0.6	24.4	5.2
Composition 3	23.8	1.6	37.1	1.0
Composition 8	4.8	1.1	6.6	0.8
Composition 4	5.9	0.9	15.5	1.0

5 Table 2

1 week after preparation	TO-stain ΔR 460		TO-stain ΔR 460	
10 FH, 2 g/l, T=25 C	2 hours after washing		1 day after washing	
Triplicate measurements	average	stdv	average	Stdv
Composition 5	11.8	1.3	13.1	
Composition 1	18.5	0.7	36.6	
Composition 6	11.2	0.4	12.8	
Composition 2	14.9	0.4	37.4	
Composition 7	13.6	0.4	18.9	
Composition 3	19.9	2.7	39.3	
Composition 8	4.1	1.0	5.5	
Composition 4	3.7	0.8	12.6	

Table 3

2 weeks after preparation	TO-stain ΔR 460		TO-stain ΔR 460	
10 °FH, 2 g/l, T=25 C	2 hours after washing		1 day after washing	
Triplicate measurements	average	stdv	average	Stdv
Composition 5	11.7	1.4	16.5	1.3
Composition 1	20.2	0.9	34.0	0.8
Composition 6	12.1	0.3	16.2	3.3
Composition 2	14.8	0.1	34.5	0.7
Composition 7	14.7	0.1	17.6	1.5
Composition 3	19.7	2.4	35.2	1.3
Composition 8	3.9	0.8	5.4	0.7
Composition 4	4.1	0.3	11.4	0.8

5 Table 4

3 weeks after preparation	TO-stain ΔR 460		TO-stain ΔR 460	
10 °FH, 2 g/l, T=25 C	2 hours after washing		1 day after washing	
Triplicate measurements	average	stdv	average	Stdv
Composition 5	13.5	0.6	16.9	2.5
Composition 1	14.1	1.7	33.8	1.0
Composition 6	12.8	0.5	17.6	3.2
Composition 2	14.5	0.5	34.1	1.4
Composition 7	16.1	1.8	18.8	4.5
Composition 3	16.6	0.9	33.9	0.3
Composition 8	3.1	0.7	4.2	1.6
Composition 4	3.9	0.8	7.8	1.2

Table 5

4 weeks after preparation	TO-stain ΔR 460		TO-stain ΔR 460	
10 °FH, 2 g/l, T=25 °C	2 hours after washing		1 day after washing	
Triplicate measurements	average	stdv	average	Stdv
Composition 5	12.1	1.1	15.0	2.2
Composition 1	17.8	1.7	34.5	1.2
Composition 6	12.3	0.8	15.2	1.9
Composition 2	16.5	1.7	34.2	1.5
Composition 7	14.1	1.9	16.7	1.6
Composition 3	14.5	0.1	28.3	1.3
Composition 8	3.5	0.6	5.0	1.0
Composition 4	3.4	2.1	9.5	1.9

5 Table 6

6 weeks after preparation	TO-stain ΔR 460		TO-stain ΔR 460	
10 °FH, 2 g/l, T=25 °C	2 hours after washing		1 day after washing	
Triplicate measurements	average	stdv	average	Stdv
Composition 5	14.8	0.9	15.8	1.5
Composition 1	18.7	1.6	34.3	1.4
Composition 6	15.2	1.0	15.5	1.3
Composition 2	16.8	0.7	31.0	1.4
Composition 7	19.1	0.7	19.8	1.5
Composition 3	16.9	0.7	17.1	0.6
Composition 8	6.2	0.7	7.1	0.5
Composition 4	6.1	0.3	7.3	0.5

Example 3

10 Composition 5 Wisk™ liquid USA, 1999

Composition 6 OMO™ liquid NL, 1999

Composition 7 OMO-liquido™ Brazil, 1999

Composition 9 non-aqueous liquid formulation:

Ingredient	Wt%
Nonionic surfactant	26.6
Monopropylene glycol	5.5
Pigment premix	0.017
Glycerol	21.36
Monoethanolamine	7.56
Oleic fatty acid	13.10
Water	Up to 100
Linear alkyl benzene sulfonate	20.1
Perfume	1.6
Protease Enzyme	1.0

In all experiments, 2 g/l of the above formulation was used, with either 2.5 or 5 microM of metal complex 1-8, or 2.5 or 5 microM of the ligand 1-8 dissolved in the wash liquor. In all cases tomato stains were used and treated further as described for Example 3. The cloths were measured immediately after drying and after 24 h storage (expressed as ΔR 460 bleaching value (a higher value indicates a cleaner cloth)).

Ligand 1: N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane (MeN4py).

Ligand 2: N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-amino-2-phenylethane (BzN4py). The synthesis of ligand 2 has been disclosed in EP 0909 809.

Ligand 3: N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-aminomethane (N4py). The synthesis of ligand 3 has been disclosed in Wo-A-9534628.

5

Ligand 4: N,N,N',N'-tetrakis(pyridin-2ylmethyl)ethane-diamine (tpen). Ligand 4 was synthesised according to a modified literature procedure (see G. Anderegg, F. Wenk, *Helv. Chim. Acta*, 50(8), 2330 (1967).

10 Trispicen-NH (5.95 g, 17.9 mmol) and 1.67 g (18.4 mmol) of 2-pyridinecarboxaldehyde were dissolved in 120 ml 1,2-dichloroethane. To this mixture $\text{NaBH}(\text{OAc})_3$ (18 mmol) was added and the mixture was refluxed for 16 h. Subsequently 50 ml of 5 N NaOH and after 1 h stirring the organic layer
15 was separated and the water layer was further extracted with dichloromethane. After drying the organic layers over sodium sulfate, filtration and evaporation of the solvents, a semi-solid paste was obtained that was purified over an alumina column (elutant: ethyl acetate/ hexane/ triethylamine
20 9:10:1). The oil isolated become now solid and could be crystallised from ethyl acetate/hexane (1/1) yielding a pale-brown powder (4.45 g, 10.5 mmol; 58.6%). ^1H -nmr (CDCl_3): δ 2.78 (s, 4H); 3.75 (s, 8H); 7.0 (m, 4H); 7.38 (m, 4H); 7.50 (m, 4H); 8.43 (m, 4H).

25

Ligand 5: N-methyl-N,N',N'-tris(3-methyl-pyridin-2ylmethyl)ethane-diamine (trilen). The synthesis of ligand 5 has been disclosed in EP 1001 009.

30 Ligand 6: N,N,N'-tris(pyridin-2ylmethyl)ethane-diamine (trispicen-NH).

First N,N'-bis(pyridin-2ylmethyl)-ethanediamine (bispicen) was synthesised by the following procedure. Ethylenediamine (26 ml, 0.38 mol) was dissolved in 200 ml dry methanol. To this mixture 74 ml (0.76 mol) pyridincarboxaldehyde was added. The mixture was refluxed for 2 h, after which the mixture was left to cool to RT and in small portions 40 g of NaBH₄ was added. The mixture was subsequently stirred for 16 h at RT. The methanol was evaporated and 500 ml of water was added. The aqueous mixture was extracted by three portions of dichloromethane (100 ml) and the dichloromethane solution was dried over sodium sulfate, filtered off and the solvent was removed. The dark oil containing N,N'-bis(pyridin-2ylmethyl)-ethanediamine (73.7 g; 81%) was analysed by NMR and used without further purification. ¹H-nmr (CDCl₃): δ 2.20 (br, NH); 2.78 (s, 4H); 3.85 (s, 4H); 7.00-7.7.40 (m, 4H); 7.58 (m, 2H); 8.45 (m, 2H).

In the second step the amination of bispicen with 2-pyridincarboxaldehyde was synthesised. 73.7 g of the unpurified bispicen material (see above) was under argon dissolved in 750 ml of dry diethylether (distilled over P₂O₅). To this solution 32.8 g of 2-pyridincarboxaldehyde was added, the reaction mixture was stirred and cooled in an ice/water bath. After 20 min a white precipitate was formed that was filtered off (P4-glass filter) and dried with dry ether. The yield was 66.6 g (66%) and was used without further purification. ¹H-nmr (CDCl₃): δ 2.75 (m, 2H); 3.13 (m, 2H); 3.65 (d, 2H); 4.93 (d, 2H); 4.23 (s, 1H); 7.00-7.90 (m, 9H); 8.43 (m, 3H).

In the third step the desired ligand was obtained (N,N,N'-tris(pyridin-2ylmethyl)ethane-diamine - trispicen-NH). The aminal (45.0 g; 0.135 mol), obtained as described as above, was dissolved in 1.2 l of dry methanol (distilled over Mg), and to this mixture 8.61 g (0.137 mol) of NaBCNH₃ was added in small portions. Subsequently 21 ml of trifluoroacetic acid was added dropwise in the solution. The mixture was stirred for 16 h at RT and subsequently 1.05 L of 5N NaOH was added and the mixture was stirred for 6 h. Extraction with dichloromethane yielded after drying, filtration and removal of the solvent a yellow oil as product (42.7 g , 0.128 mol; 95%. ¹H-nmr (CDCl₃): δ 2.15 (br, NH); 2.75 (s, 4H); 3.80 (s, 4H); 3.82 (s, 2H); 7.0-7.8 (m, 3H); 7.45-7.70 (m, 6H); 8.40-8.60 (m, 3H). ¹³C-nmr (CDCl₃): δ 53.9 (t); 54.7 (t); 60.4 (t); 121.7 (d); 121.9 (d); 122.1 (d); 123.0 (d); 136.3 (d); 136.4 (d); 148.9 (d); 149.1 (d); 159.3 (s); 159.6 (s).

Ligand 7: N-methyl-,N,N'N'-tris(pyridin-2ylmethyl)ethane-diamine (trispicen-NMe). Ligand 7 was prepared according to a modified procedure described by Bernal et al (J. Chem. Soc., Dalton Trans, 22, 3667 (1995)).

Trispicen-NH (10g, 30 mmol) was dissolved in 25 ml formic acid and 10 ml water. To this mixture 36 % formaldehyde solution was added (16 ml, 90 mmol) and the mixture was warmed up till 90 °C for 3 h. Formic acid was evaporated and the 2.5 N NaOH solution was added until the pH was higher than 9. Extraction by dichloromethane and drying over sodium sulfate, filtration of the solution and subsequently drying yielded a dark-coloured oil (8.85g). The oil was purified over a alumina column (elutant: ethyl acetate/ hexane/

triethylamine 9:10:1). Yield 7,05g pale yellow oil (20,3mmoles; 68%). ^1H -nmr (CDCl_3): δ 2.18 (s, 3H); 2.65 (m, 2H); 2.75 (m, 2H); 3.60 (s; 2H); 3.83 (s; 4H); 7.10 (m, 3H); 7.3-7.6 (m, 6H); 8.5 (d, 3H).

5

Ligand 8: tris(pyridin-2-ylmethyl)amine (tpa)

Ligand 8 was prepared according to literature procedures (see G. Anderegg, F. Wenk,, *Helv. Chim. Acta*, 50(8), 2330 (1967)).

10

Complex 1: $[(\text{MeN4Py})\text{FeCl}]\text{Cl}$

The synthesis of Complex 1 is described in Example 1.

Complex 2: $[(\text{BzN4Py})\text{Fe}(\text{CH}_3\text{CN})](\text{ClO}_4)_2$

15 The synthesis of Complex 2 is described in EP 0909 809. An optimised synthetic procedure is given below: 3.0 g (6.56 mmol) of BzN4Py was dissolved in 30ml methanol and 30ml acetonitrile. 2.26 g (6.23 mmol) of $\text{Fe}(\text{ClO}_4) \cdot 6\text{H}_2\text{O}$ (Aldrich) was added to solution containing the ligand in small
20 portions. To the dark-red coloured solution in total 100 ml of ethyl acetate was added to facilitate the crystallisation procedure. After 18 h stirring, the red powder was filtered off, washed with ethyl acetate and dried, yielding 3.85 g of the desired complex (anal: see EP 0909 809).

25

Complex 3: $[(\text{N4Py})\text{FeCl}]\text{Cl}$

Complex 3 was synthesised according to the procedure as described for the analogous MeN4py complex using now N4py as ligand (see example 1).

30

Complex 4: $[(\text{tpen})\text{Fe}](\text{ClO}_4)_2$

Complex 4 was prepared according to the procedure found in H. Toftlund et al., J.Am. Chem. Soc., 112, 6814 (1990)

Complex 5: $[(\text{trilen})\text{FeCl}]\text{PF}_6$

5 Complex 5 was prepared according to EP 1001 009

Complex 6: $[(\text{trispicen-NH})\text{FeCl}]\text{PF}_6$

Trispicen-NH (8.0 g; 24.0 mmol) was dissolved in 60 ml methanol/water 1/1 v/v) and was heated till 50 °C. $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ 10 4,78g; 24,0mmoles) was added in small portions. The dark blue-purple solution was stirred for 10 min at 50 °C. Subsequently 4.42 g (24 mmol) of KPF_6 was added and the solution was stirred for 2 days at RT. The dark powder was filtered, washed with methanol/water and then with ethyl 15 acetate. The powder was dried in the air. Yield 11.6 g.

Complex 7: $[(\text{trispicen-NMe})\text{FeCl}]\text{PF}_6$

TrispicenNMe (6,0g; 17,3mmoles) was dissolved in 15 ml methanol/water 1/1 v/v) and was heated till 50 °C. $\text{FeCl}_2 \cdot 4\text{H}_2\text{O}$ 20 3,43g; 17,0mmoles), dissolved in 20 ml water/methanol 1/1), was added. The dark solution was stirred for 20 min at 50 °C. Subsequently 3.17 g (17 mmol) of KPF_6 dissolved in 10 ml water, was added and the solution was stirred for 15 h to yield a yellow precipitation. The solid was filtered off, 25 wasged with methanol/water 1/1, v/v) and ethyl acetate. Drying yielded 8.25 g of a pale-yellow powder.

Complex 8: $[\text{Fe}_2(\text{tpa})_2(\text{H}_2\text{O})_2](\text{ClO}_4)_2$

Complex 8 was kindly donated by Prof. L. Que, University of 30 Minnesota, USA (references: L. Que et al., Inorg Chim. Acta,

273, 393 (1998) and H. Toftlund et al., Inorg. Chem., 33, 3127 (1994).

Table 7. Bleaching results obtained on tomato stains for the
5 different complexes (5 microM) in solutions containing the
four liquid formulations (compositions 5, 6, 7 and 9). The
bleaching results obtained immediately after drying (t=0)
and after 1 day storage are shown. All values expressed in
 ΔR 460 values; typical errors are in the order of 2 points.

10

	Comp 5		Comp 6		Comp 7		Comp 9	
	t=0	t=1	t=0	t=1	t=0	t=1	t=0	t=1
Complex 1	20	50	41	47	35	55	42	49
Complex 2	20	48	42	50	31	51	42	52
Complex 3	31	49	35	50	31	53	44	52
Complex 4	16	39	16	23	26	48	29	42
Complex 5	33	47	36	46	39	52	43	50
Complex 6	15	22	12	15	16	23	15	18
Complex 7	19	39	17	20	25	46	27	33
Blank	11	13	15	19	13	14	15	18

From these results is clear that especially complexes 1, 2,
3, and 5 give a good tomato stain bleaching with air,
15 although the exact amount depends on the formulation
employed. Complexes 4, 6, and 7 give somewhat lower
bleaching activity, but still in most cases more than the
blanks.

Table 8. Bleaching results obtained on tomato stains for the different ligands (5 microM) in solutions containing the four liquid formulations (compositions 5, 6, 7 and 9). The bleaching results obtained immediately after drying (t=0) and after 1 day storage are shown. All values expressed in ΔR 460 values; typical errors are in the order of 2 points.

	Comp 5		Comp 6		Comp 7		Comp 9	
	t=0	t=1	t=0	t=1	t=0	t=1	t=0	t=1
Ligand 1	16	42	22	44	18	32	33	52
Ligand 2	16	40	26	47	16	34	32	51
Ligand 3	18	37	19	39	18	39	33	53
Ligand 5	22	40	26	36	19	36	41	52
Ligand 6	14	16	14	16	14	20	18	20
Ligand 7	16	20	16	19	19	28	19	22
Blank	11	13	14	19	12	14	15	18

All ligands in the wash liquor containing the four formulations give significant enhancement of the tomato stain bleaching in the air. This effect is especially clear for ligands 1, 2, 3 and 5.

Table 9. Bleaching results obtained on tomato stains for the different complexes (2.5 microM) in solutions containing the four liquid formulations (compositions 5, 6, 7 and 9). The bleaching results obtained immediately after drying (t=0) and after 1 day storage are shown. All values expressed in ΔR 460 values; typical errors are in the order of 2 points.

	Comp 5		Comp 6		Comp 7		Comp 9	
	t=0	t=1	t=0	t=1	t=0	t=1	t=0	t=1
Complex 1	15	46	38	48	27	49	22	44
Complex 2	22	46	15	35	28	47	18	38
Complex 3	20	46	24	43	27	44	24	47
Complex 4	12	19	9	11	18	24	12	19
Complex 5	30	43	23	33	27	36	23	40
Complex 6	9	10	8	9	16	23	13	14
Complex 7	15	18	9	10	23	32	17	21
Complex 8	10	13	11	12	11	13	12	15
Blank	10	11	9	10	12	14	11	12

Table 10. Bleaching results obtained on tomato stains for the different ligands (2.5 microM) in solutions containing the four liquid formulations (compositions 5, 6, 7 and 9).

- 5 The bleaching results obtained immediately after drying (t=0) and after 1 day storage are shown. All values expressed in ΔR 460 values; typical errors are in the order of 2 points.

	Comp 5		Comp 6		Comp 7		Comp 9	
	t=0	t=1	t=0	t=1	t=0	t=1	t=0	t=1
Ligand 1	13	26	9	13	11	13	11	15
Ligand 2	11	19	10	14	10	13	13	21
Ligand 3	13	26	9	11	12	14	13	17
Ligand 5	13	20	9	11	12	16	14	19
Ligand 6	11	12	10	12	10	11	10	12
Ligand 7	12	15	9	11	10	12	13	15
Ligand 8	8	9	9	11	13	15	11	14
Blank	10	11	9	10	11	14	11	12

Discussion of Results:

The results show that the activity of FeMeN4PyCl_2 is stable for six weeks in the detergent Compositions 1 and 2.

However, the activity of FeMeN4PyCl_2 in composition 4 and in
5 composition 3 after more than four weeks storage decreased.

Without being bound by theory, it is more than likely that STP present in a liquid composition gives the negative effect on the storage stability and that addition of iron salt restores the activity. The results show that by adding

10 a liquid composition containing a ligand or transition metal complex thereof to the wash liquor a bleaching capacity is provided without the presence of an added peroxy species or precursor thereof. In addition, the bleaching capacity is provided at a low concentration of a ligand or transition
15 metal complex thereof in the wash liquor.

1) FeMeN4PyCl_2 , amongst others, gives clear bleach benefits in a variety of liquid formulations (incl. rinse conditioner) on tomato-oil stains.

20 2) The bleach effect upon 24 hr storage of the cloths in the dark is much larger then 2 h after the wash.

3) No visual change in structural phase after two weeks.

4) Immediate colour change upon addition of FeMeN4PyCl_2 of the liquid observed.

25 5) Similar bleach performance upon 6 weeks of storage as found immediately after mixing for the detergent Compositions 1 and 2, implying a stable system.

6) No bleach effects were more observed after 6 weeks of storage for detergent Composition 3 and rinse conditioner

30 Composition 4.

Complex 8 and ligand 8 show significant decreased bleach benefit in a liquid bleach composition. As is known from inorganic chemistry, in general pentadentate ligands give rise to more stable complexes than tetradentate ligands; this is known as the chelate effect. (see Huheey, inorganic chemistry, 2nd edition, Harper and Row). The decreased stability is especially noted in basic aqueous media, where formation of insoluble iron Hydroxide species are often encountered. The decreased stability of the iron tpa complexes/species gives rise to a poorer performance in the liquid detergent formulations.

There are many liquid formulations for detergents and rinse conditioners or other liquid products that may be enhanced by conferring a bleaching ability to the liquid formulation. As will be evident to one skilled in the art the present invention is applicable to known liquid formulations and liquid formulations to be developed.

As one skilled in the art will appreciate determining the suitability of a particular catalyst for bleaching of a substrate by atmospheric oxygen in a particular liquid formulation is a matter of routine experimentation. The present invention extends to both isotropic and complex liquid compositions and formulations a brief discussion of which follows. Some isotropic formulations are termed 'micro-emulsion' liquids that are clear and thermodynamically stable over a specified temperature range. The 'micro-emulsion' formulation may be water in oil, or oil in water emulsions. Some liquid formulations are macro-emulsions that are not clear and isotropic. Emulsions are

considered meta-stable. Concentrated, clear compositions containing fabric softening actives have been disclosed in WO 98/08924 and WO 98/4799, both Procter & Gamble. Such compositions comprise bio-degradable fabric conditioners.

5 However, both disclose compositions comprising water miscible solvents that do not form water-in-oil micro-emulsions. Clear fabric conditioning compositions have also been disclosed in EP 730023 (Colgate Palmolive), WO 96/19552 (Colgate Palmolive), WO 96/33800 (Witco Co.), WO 97/03170
10 (Procter & Gamble), WO 97/03172 (Procter & Gamble), WO 97/03169 (Procter & Gamble), US 5492636 (Quest Int.) and US 5427697 (Procter & Gamble). Liquid formulations of the present invention may contain for example; monoethoxy quats; AQAs and bis-AQAs; cationic amides; cationic esters;
15 amino/diamino quats; glucamide; amine oxides; ethoxylated polyethyleneimines; enhancement polymers of the form linear amine based polymers, e.g. bis-hexamethylenetriamine; polyamines e.g. TETA, TEPA or PEI polymers.

20 Experimentation to determine catalyst-liquid stability, as detailed above, may be varied. The aforementioned method determined the catalyst-liquid stability/compatibility by examining how the oxygen bleaching ability of a particular catalyst-liquid formulation varied with time.

25 Alternatively, the determination may be conducted by monitoring the concentration of a particular catalyst in a liquid formulation by known techniques, for example NMR, HPLC, Liquid Chromatography-Mass Spectroscopy, Infra Red, UV-visible measurements, etc, over a period of time.

30 Alternatively, another possible method of determining catalyst-liquid stability would be to analyse the activity

- 91 -

of a certain transition metal compound by oxidation activity studies using a dye/compound that gives a colour change upon oxidation. An example of a dye/compound that gives a colour change upon oxidation is 2,2'-azinobis(3-


5 ethylbenzothiazoline-6-sulfonate) and many other
dyes/compounds that give a colour change upon oxidation are
known. Methods for using a dye/compound that gives a colour
change upon oxidation are known in the art for establishing
activity of a variety of redox enzymes.

10

CLAIMS:

1. A liquid bleaching composition comprising an organic substance which forms a complex with a transition metal, the complex catalysing bleaching of a substrate by atmospheric oxygen, and a liquid carrier or solvent, wherein the composition is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system.

2. A liquid bleaching composition according to claim 1, wherein the organic substance comprises a pentadentate ligand of the general formula (B):



(B)

wherein

each R^1 , R^2 independently represents $-R^4-R^5$,

R^3 represents hydrogen, optionally substituted alkyl, aryl or arylalkyl, or $-R^4-R^5$,

each R^4 independently represents a single bond or optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene, alkylene ether, carboxylic ester or carboxylic amide, and

each R^5 independently represents an optionally N-substituted aminoalkyl group or an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl,

pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl,
pyrimidinyl, triazolyl and thiazolyl.

3. A liquid bleaching composition according to claim 2,
5 wherein the ligand is N,N-bis(pyridin-2-yl-methyl)-1,1-
bis(pyridin-2-yl)-1-aminoethane.

4. A liquid bleaching composition according to claim 1,
wherein the medium has a pH value in the range from pH 6 to
10 11.

5. A liquid bleaching composition according to claim 4,
wherein the medium has a pH value in the range from pH 7 to
10 10.

6. A liquid bleaching composition according to claim 4,
wherein the medium is substantially devoid of a transition
metal sequestrant.

7. A liquid bleaching composition according to claim 6,
wherein the medium further comprises a surfactant.

8. A liquid bleaching composition according to claim 4,
wherein the medium further comprises a builder.

9. A liquid bleaching composition according to claim 1,
wherein the organic substance comprises a preformed complex
of a ligand and a transition metal.

- 94 -


10. A liquid bleaching composition according to claim 1, wherein the organic substance comprises a free ligand that complexes with a transition metal present in the water.

5 11. A liquid bleaching composition according to claim 1, wherein the organic substance comprises a free ligand that complexes with a transition metal present in the substrate.

10 12. A liquid bleaching composition according to claim 1, wherein the organic substance comprises a composition of a free ligand or a transition metal-substitutable metal-ligand complex, and a source of transition metal.

15 13. A method of bleaching a substrate comprising applying to the substrate a liquid bleaching composition that comprises an organic substance which forms a complex with a transition metal, the complex catalysing bleaching of the substrate by atmospheric oxygen, and a liquid carrier or solvent, wherein the composition is substantially devoid of
20 peroxygen bleach or a peroxy-based or -generating bleach system.

14. A method to claim 13, wherein the organic substance comprises a pentadentate ligand of the general formula (B):



(B)

wherein

each R^1 , R^2 independently represents $-R^4-R^5$,

R^3 represents hydrogen, optionally substituted alkyl, aryl or arylalkyl, or $-R^4-R^5$,

5 each R^4 independently represents a single bond or optionally substituted alkylene, alkenylene, oxyalkylene, aminoalkylene, alkylene ether, carboxylic ester or carboxylic amide, and

each R^5 independently represents an optionally N-
10 substituted aminoalkyl group or an optionally substituted heteroaryl group selected from pyridinyl, pyrazinyl, pyrazolyl, pyrrolyl, imidazolyl, benzimidazolyl, pyrimidinyl, triazolyl and thiazolyl.

15 15. A method according to claim 14, wherein the ligand is N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane.

20 16. A method according to claim 14, wherein the method is conducted in a medium having a pH value in the range from pH 6 to 11.

25 17. Use of an organic substance which forms a complex with a transition metal, the complex catalysing bleaching of a substrate by the atmospheric oxygen, as a catalytic bleaching agent in a liquid bleaching composition substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system.

30 18. A method of treating a textile by contacting the textile with a liquid bleaching composition that comprises

an organic substance which forms a complex with a transition metal, the complex catalysing bleaching by atmospheric oxygen, and a liquid carrier or solvent, wherein the composition is substantially devoid of peroxygen bleach or a peroxy-based or -generating bleach system, whereby the complex catalyses bleaching of the textile by atmospheric oxygen after the treatment.

19. A liquid bleaching composition according to claim 1, wherein the organic substance comprises a pentadentate ligand.

20. A liquid bleaching composition according to claim 19, wherein the pentadentate ligand is in the form of an iron complex.

21. A liquid bleaching composition according to claim 19 having a pH value in the range from pH 7 to 10 comprising N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane, the composition substantially devoid of a transition metal sequestrant having a higher binding affinity for iron ions than N,N-bis(pyridin-2-yl-methyl)-1,1-bis(pyridin-2-yl)-1-aminoethane.

ABSTRACT:**COMPOSITION AND METHOD FOR BLEACHING A SUBSTRATE**

5 The invention relates to a liquid bleaching composition for
catalytically bleaching substrates, especially laundry
fabrics, with atmospheric oxygen or air. A liquid bleaching
composition is provided comprising an organic substance
which forms a complex with a transition metal, the complex
10 catalysing bleaching of a substrate by atmospheric oxygen,
and a liquid carrier or solvent, wherein the composition is
substantially devoid of peroxygen bleach or a peroxy-based
or -generating bleach system. Also provided is a method of
bleaching a substrate comprising applying the liquid
15 bleaching composition to the substrate. Also provided is a
method of treating a textile by contacting the textile with
the liquid bleaching composition, whereby the complex
catalyses bleaching of the textile by atmospheric oxygen
after the treatment.

20

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY
(Includes Reference to PCT International Applications)ATTORNEY'S DOCKET NUMBER
C4007 (C)

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

COMPOSITION AND METHOD FOR BLEACHING A SUSTRATE

the specification of which (check only one item below):

☒ is attached hereto.☐ was filed as United States application Serial No. _____ on _____
and was amended on _____ (if applicable)☐ was filed as PCT International application _____ on _____
and was amended under PCT Article 19 on _____ (if applicable)

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby claim foreign priority benefits under Title 35, United States Code §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:

COUNTRY (if PCT indicate PCT)	APPLICATION NUMBER	DATE OF FILING (day, month, year)	PRIORITY CLAIMED UNDER 35 USC 119	
P.C.T.	PCT/GB99/02876	1 ST September 1999	<input checked="" type="checkbox"/> YES	<input type="checkbox"/> NO
P.C.T.	PCT/GB99/02878	1 ST September 1999	<input checked="" type="checkbox"/> YES	<input type="checkbox"/> NO
UNITED KINGDOM	0004990.8	1 st March 2000	<input checked="" type="checkbox"/> YES	<input type="checkbox"/> NO
			<input type="checkbox"/> YES	<input type="checkbox"/> NO

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that /those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code §112. I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations §1.56 (a) which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application.

PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT UNDER 35 U.S.C. 120.

U.S. APPLICATION(S)		STATUS (CHECK ONE)		
U.S. APPLICATION NUMBER	U.S. Filing Date	PATENTED	PENDING	ABANDONED
PCT APPLICATIONS DESIGNATING THE U.S.				
PCT APPLICATION NUMBER	PCT Filing Date	U.S. Serial Numbers Assigned (if any)		

COMBINED DECLARATION FOR PATENT APPLICATION AND POWER OF ATTORNEY (continued)
(includes Reference to PCT International Applications)

ATTORNEY DOCKET NUMBER
C4007 (C)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

CUSTOMER NUMBER: 000201

Direct all correspondence to: CUSTOMER NUMBER 000201

201

Full Name of Inventor	Family Name HAGE	First Given Name Ronald	Second Given Name
Residence & Citizenship	City 3133 AT Vlaardingen	State or Foreign Country Netherlands	Country of Citizenship Netherlands
Post Office Address	Post Office Address c/o Unilever Research Vlaardingen Olivier van Noortlaan 120	City 3133 AT Vlaardingen	State & Zip Code/Country Netherlands

202

Full Name of Inventor	Family Name VEERMAN	First Given Name Simon	Second Given Name Marinus
Residence & Citizenship	City 3133 AT Vlaardingen	State or Foreign Country Netherlands	Country of Citizenship Netherlands
Post Office Address	Post Office Address c/o Unilever Research Vlaardingen Olivier van Noortlaan 120	City 3133 AT Vlaardingen	State & Zip Code/Country Netherlands

203

Full Name of Inventor	Family Name	First Given Name	Second Given Name
Residence & Citizenship	City	State or Foreign Country Netherlands	Country of Citizenship
Post Office Address	Post Office Address	City	State & Zip Code/Country

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such wilful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor 201	Signature of Inventor 202	Signature of Inventor 203
Date	Date	Date